

Warm water is mixing up life in the Arctic

Unique 15-year observation series reveals how sensitive marine ecosystems in polar regions are to change

Summary: The warming of arctic waters in the wake of climate change is likely to produce radical changes in the marine habitats of the High North. This is indicated by data from long-term observations in the Fram Strait, which researchers from the Alfred Wegener Institute (AWI) have now analysed. Their most important finding: even a short-term influx of warm water into the Arctic Ocean would suffice to fundamentally impact the local symbiotic communities, from the water's surface down to the deep seas. The study reports what happened between 2005 and 2008 in *Ecological Indicators*.


The Arctic is a remote and extreme habitat. Yet, despite its isolation, the anthropogenically accelerated climate change will massively transform this region in the years to come. We've known for some time now that, thanks to the greenhouse effect, the sea-ice cover in the Arctic is shrinking and the world's oceans are gradually warming. But it was unclear just how polar marine organisms were responding.

Scientists at the Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research (AWI) have now used an unprecedented set of long-term observations to show that arctic marine habitats could radically change if subjected to a sustained rise in temperature. The most surprising finding is that the thermally induced changes at the ocean surface can rapidly spread to affect life in the deep seas.

In order to investigate the changes in the ecosystems of arctic waters, for the past 15 years the AWI has operated the deep-sea observatory "HAUSGARTEN" in the Fram Strait, the sea lane between Greenland and Svalbard. HAUSGARTEN is a network of 21 individual research stations, which AWI staff seeks out every summer to collect water and soil samples. Some of the stations feature anchored systems that operate year-round, recording the water temperature and tides, collecting water and soil samples at regular intervals, and capturing the sediments that drift down to the seafloor from the upper water layers.

This is the only observatory of its kind in the world. There's no other project in which readings from the surface down to the ocean floor were taken in fixed positions over such a long time -- let alone in the polar regions, says AWI biologist Thomas Soltwedel.

For the current publication, the AWI researcher and his team analysed the first 15 years of the HAUSGARTEN dataset. The Fram Strait is especially interesting for Soltwedel and his colleagues because it represents the only deep juncture in the Arctic Ocean, allowing water masses from the Atlantic to flow into the Arctic to the west of Svalbard (see figure). In turn, water and ice floes find their way back out of the Arctic Ocean on the strait's Greenland side.



Normally the water near the surface, which flows north out of the Atlantic through the Fram Strait, has an average temperature of 3°C. Yet with the help of their observatory, the AWI researchers were able to determine that from 2005 to 2008 the average temperature of the inflowing water was one to two degrees higher: In that time, large quantities of warmer water poured into the Arctic Ocean. Since polar organisms have adapted to living in constant cold, this extra heat input hit them like a temperature shock, explains Soltwedel.

The reactions in the ecosystem were correspondingly extreme; the researchers were

able to identify serious changes in various symbiotic communities, from microorganisms and algae to zooplankton. One major change was the increase in free-swimming conchs and amphipods, which are normally found in the more temperate and subpolar regions of the Atlantic. In contrast, the number of conchs and amphipods in the Arctic dropped significantly.

Another remarkable aspect is the decline in small, hard-shelled diatoms. Prior to the unexpected influx of warm water, they made up roughly 70% of the vegetable plankton in the Fram Strait. But during the warm phase, the foam algae *Phaeocystis* took their place. Unlike diatoms, foam algae tend to clump and sink to the ocean floor, where they become a food source. But the sudden rise in available food led to major changes in deep-sea life, including a noticeable increase in the settlement density of benthic organisms, explains Soltwedel.

The study is not able to exactly conclude how all these changes will affect the overall arctic food web in the future.

Since the flow of warm water has subsided, the water temperature in the Fram Strait has stabilized -- though it is still slightly above the average value from before 2005. Yet the changes in the ecosystem have partly become lasting realities. For example, the number of diatoms remains very small. And the conchs from the lower latitudes seem to have made a home for themselves in the Fram Strait.


At the moment they can't say whether the warm-water influx they have monitored is due to climate change, as there are also natural climate fluctuations that occur every ten years. They overlap with the anthropogenically caused effects of climate change, as a result of which, after 15 years, the researchers don't know whether this flow of warm water was the result of strictly natural causes. However, the results of the ecological long-term studies clearly show that even short-term changes in ocean temperature can drastically impact life in the Arctic.

Without the observations made over the years at HAUSGARTEN, which are supplemented by regular ship-borne measurements, that glimpse would have never been possible. This study indicates how important long-term observations at the same location are necessary to identify lasting changes and making forecasts on future developments.

[pdf reprint follows.....](#)



Contents lists available at ScienceDirect



Ecological Indicators

journal homepage: www.elsevier.com/locate/ecolind

Natural variability or anthropogenically-induced variation? Insights from 15 years of multidisciplinary observations at the arctic marine LTER site HAUSGARTEN

Thomas Soltwedel^{a,*}, Eduard Bauerfeind^a, Melanie Bergmann^a, Astrid Bracher^{a,b}, Nataliya Budaeva^{c,d}, Kathrin Busch^a, Alexandra Cherkasheva^{b,e}, Kirsten Fahl^a, Katarzyna Grzelak^f, Christiane Hasemann^a, Marianne Jacob^a, Angelina Kraft^g, Catherine Lalande^h, Katja Metfies^a, Eva-Maria Nöthig^a, Kirstin Meyerⁱ, Nadia-Valérie Quéric^j, Ingo Schewe^a, Maria Włodarska-Kowalcuk^f, Michael Klages^k

^a Alfred Wegener Institute, Helmholtz-Centre for Polar and Marine Research, Am Handelshafen 12, 27570 Bremerhaven, Germany

^b University of Bremen, Institute of Environmental Physics, PO Box 330440, 28334 Bremen, Germany

^c P.P. Shirshov Institute of Oceanology, Russian Academy of Sciences, 36, Nakhimovsky Pr., Moscow 117997, Russia

^d Natural History Collections, University Museum of Bergen, Allégaten 41, 5020 Bergen, Norway

^e St Petersburg State University, 7-9, Universitetskaya nab., St. Petersburg 199034, Russia

^f Institute of Oceanology, Polish Academy of Sciences, 81-712 Sopot, Powstańców Warszawy 55, Poland

^g German National Library of Science and Technology TIB, Welfengarten 1B, 30167 Hannover, Germany

^h Québec-Océan, Département de biologie, Université Laval, Québec, Canada G1V 0A6

ⁱ Oregon Institute of Marine Biology, 63466 Boat Basin Road, Charleston, OR 97420, USA

^j Department of Geobiology, Center of Earth Sciences, University of Göttingen, Goldschmidtstrasse 3, 37077 Göttingen, Germany

^k Sven Lovén Centre for Marine Sciences at the University of Gothenburg, Kristineberg 566, 451 78 Fiskebäckskil, Sweden

ARTICLE INFO

Article history:

Received 29 April 2015

Received in revised form

29 September 2015

Accepted 2 October 2015

Keywords:

HAUSGARTEN

Arctic Ocean

Deep sea

Natural variability

Anthropogenic impact

ABSTRACT

Time-series studies of arctic marine ecosystems are rare. This is not surprising since polar regions are largely only accessible by means of expensive modern infrastructure and instrumentation. In 1999, the Alfred Wegener Institute, Helmholtz-Centre for Polar and Marine Research (AWI) established the LTER (Long-Term Ecological Research) observatory HAUSGARTEN crossing the Fram Strait at about 79° N. Multidisciplinary investigations covering all parts of the open-ocean ecosystem are carried out at a total of 21 permanent sampling sites in water depths ranging between 250 and 5500 m. From the outset, repeated sampling in the water column and at the deep seafloor during regular expeditions in summer months was complemented by continuous year-round sampling and sensing using autonomous instruments in anchored devices (i.e., moorings and free-falling systems). The central HAUSGARTEN station at 2500 m water depth in the eastern Fram Strait serves as an experimental area for unique biological in situ experiments at the seafloor, simulating various scenarios in changing environmental settings. Long-term ecological research at the HAUSGARTEN observatory revealed a number of interesting temporal trends in numerous biological variables from the pelagic system to the deep seafloor. Contrary to common intuition, the entire ecosystem responded exceptionally fast to environmental changes in the upper water column. Major variations were associated with a Warm-Water-Anomaly evident in surface waters in eastern parts of the Fram Strait between 2005 and 2008. However, even after 15 years of intense time-series work at HAUSGARTEN, we cannot yet predict with complete certainty whether these trends indicate lasting alterations due to anthropologically-induced global environmental changes of the system, or whether they reflect natural variability on multiyear time-scales, for example, in relation to decadal oscillatory atmospheric processes.

© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author. Tel.: +49 471 4831 1775; fax: +49 471 4831 1776.

E-mail address: Thomas.Soltwedel@awi.de (T. Soltwedel).

1. Introduction

While always fluctuating, the global climate is presently experiencing a period of constantly increasing temperatures, with a warming trend amplified in the Arctic (Hassol, 2004). Results of large-scale simulations of the Earth's future climate by several global climate models predict a continuous increase in air and water temperatures, also leading to further reduction in ice-cover (IPCC, 2013). Since the 1950s, sea-ice retreat in the Arctic Ocean has been relatively modest at rates of 3–4% per decade (Parkinson et al., 1999). However, since the late 1990s, annual-averaged shrinking rates accelerated to 10.7% per decade (Comiso et al., 2008), whilst the summer sea-ice extent has shrunk even more rapidly. According to the US National Snow and Ice Data Center (NSIDC), arctic sea-ice during the 2012 melt season has reached its lowest extent since satellites began measuring sea-ice in 1979, with 44% ice coverage below the 1981–2010 average, and 16% ice coverage below the previous minimum extent in 2007.

Moreover, according to the results from large-scale surveys carried out by US submarines equipped with upward looking sonar and ICESat satellite observations, there has been a significant thinning of the sea-ice by approximately 50% since the late 1950s (Kwok and Rothrock, 2009). Laxon et al. (2013) compared satellite and model-derived sea-ice volume data between the years 2003–2008 and 2010–2012, and found that sea-ice volume declined by 36.2% at the end of summer, and by 9.1% at the end of winter. In its recent report, the Intergovernmental Panel on Climate Change (IPCC) prophesied that the Arctic could become ice-free in the second half of this century, while Wang and Overland (2012) argued that this scenario might even take place much earlier, with predictions as early as the end of the arctic summer 2040. Greenhouse gases emitted through human activities and the resulting increase in global mean temperatures are most likely the underlying causes of the sea-ice decline, although it is recognized that the current sea-ice decline most likely resulted from a complex interaction between natural and anthropogenic causes and variations in climate change, acting on different time scales (Drinkwater et al., 2014).

The shift from an ice-covered and cold ocean to an ice-free and warmer ocean will have severe impacts on the polar marine ecosystem and its functioning (e.g., Wassmann et al., 2011). Thinner ice may permit better growth of ice algae, but earlier and faster spring melting may reduce their growing season (Arrigo, 2013). Locally, thinner ice floes with a multitude of melt ponds allow enhanced light transmission and may increase phytoplankton growth beneath the ice (Arrigo et al., 2012; Nicolaus et al., 2012). The timing and location of pelagic primary production will generally be altered (Kahru et al., 2010). Ice-edge phytoplankton blooms will be displaced progressively northwards (Engelsen et al., 2002). Whether sea-ice retreat generally leads to an increase in primary productivity is under debate (Arrigo et al., 2008; Tremblay and Gagnon, 2009; Wassmann et al., 2010). In fact, biogeochemical models predict no or even negative changes in productivity and export flux in the Barents Sea and the Fram Strait (Forest et al., 2010; Slagstad et al., 2011).

Altered algal abundance and composition will affect zooplankton community structure (Caron and Hutchins, 2013) and subsequently the flux of particulate organic matter to the seafloor (Wohlers et al., 2009), where the changing quantity and quality of this matter will impact benthic communities (Kortsch et al., 2012; Jones et al., 2013). Changes in the predominance of certain trophic pathways will have cascading effects propagating through the entire marine community.

Generally, arctic marine organisms will be compromised by temperature regimes approaching the limits of their thermal capacity (Burrows et al., 2011, 2014). As a consequence, warmer waters in the Arctic will allow a northward expansion of sub-arctic and

boreal species (Hirche and Kosobokova, 2007; Poloczanska et al., 2013). Besides water temperature increase, expanding ocean acidification will pose another threat to pelagic and benthic life in the Arctic Ocean (e.g., Bates et al., 2009; Lischka and Riebesell, 2012; AMAP, 2013).

To detect and track the impact of large-scale environmental changes on the marine ecosystem in the transition zone between the northern North Atlantic and the central Arctic Ocean, the Alfred Wegener Institute, Helmholtz-Centre for Polar and Marine Research (AWI) established the LTER (Long-Term Ecological Research) observatory HAUSGARTEN in the Fram Strait between NE Greenland and the Svalbard archipelago (Soltwedel et al., 2005). Since 1999, repeated sampling in the water column and at the seafloor during yearly expeditions in summer months was complemented by continuous year-round sampling and sensing using autonomous instruments on anchored devices. The central HAUSGARTEN station at about 79° N, 04° E in the eastern Fram Strait (~2500 m water depth) serves as an experimental area for unique biological experiments at the deep seafloor, simulating various scenarios in changing environmental settings (Premke et al., 2006; Gallucci et al., 2008; Kanzog et al., 2009; Guilini et al., 2011; Soltwedel et al., 2013).

Time-series studies at the HAUSGARTEN observatory provide insights into processes and dynamics within an arctic marine ecosystem and act as a baseline for further investigations of ongoing changes in the Fram Strait. Long-term observations at HAUSGARTEN significantly contribute to the global community's efforts to understand variations in ecosystem structure and functioning on seasonal to decadal time-scales in an overall warming Arctic and will allow for improved future predictions under different climate scenarios.

2. Site description

The LTER observatory HAUSGARTEN is located in a region which is influenced by the highly productive Marginal Ice Zone (MIZ) in the Fram Strait. Currently, HAUSGARTEN constitutes a network of 21 permanent sampling sites, the majority of which are located along a bathymetric transect between ~250 m and ~5500 m water depth at about 79° N from the Kongsfjorden (Svalbard) in the east, along the Vestnesa Ridge toward the Molloy Hole (i.e., the deepest known depression in the Arctic Ocean) and across the Greenland continental margin (stations in the western Fram Strait were newly established in 2014). Three sampling sites close to the ice edge between 79°30' N and 80°00' N in the north-eastern Fram Strait and a supplementary site in a permanently ice-free area at 78°30' N in the eastern part of the strait complete the network (Fig. 1).

The Fram Strait is the only deep water connection between the Nordic Seas and the central Arctic Ocean with a sill depth of ca. 2600 m. The hydrography in the eastern part of the strait is characterized by the inflow of relatively warm and nutrient-rich Atlantic Water (AW) into the central Arctic Ocean (Beszczynska-Möller et al., 2012). Cooler and less-saline Polar Water exits the central Arctic Ocean as the Eastern Greenland Current (EGC) in the western part of the Fram Strait (de Steur et al., 2009), separated by a frontal system (East Greenland Polar Front) from the water masses in the eastern part of the Fram Strait (Paquette et al., 2012). Hydrographic patterns in the strait result in a variable sea-ice cover, with predominantly ice-covered areas in the west, permanently ice-free areas in the south-east, and seasonally-varying ice conditions in the central and north-eastern parts. Anomalies in summer sea-ice coverage from a 30-year reference period (1981–2000) in the greater Fram Strait since the establishment of HAUSGARTEN observatory in 1999 are shown in Fig. 2. Strong positive values (more sea-ice compared to the long-term mean) were found in 2003, 2008, and 2009,




Fig. 1. Sampling sites (●) of the LTER observatory HAUSGARTEN, geographic features, and major currents in the Fram Strait (KF: Kongsfjorden; VR: Vestnesa Ridge; MH: Molloy Hole; GCM: Greenland Continental Margin; WSC: West Spitsbergen Current; EGC: Eastern Greenland Current; RAC: Return Atlantic Current; YB: Yermak Branch, and SB: Svalbard Branch of the WSC).




Fig. 2. Sea-ice anomalies from a 30-year reference period (1981–2000) in the greater Fram Strait for 1999 through 2014 (provided by the National Snow & Ice Data center, NSIDC); the trapezoid depicts the HAUSGARTEN area.

whereas negative values (less ice compared to the long-term mean) occurred in 2001 and 2002, in 2004 through 2006, and in 2010; all other years showed intermediate values for the ice coverage at the observatory. In this way, the data showed no clear temporal trend in sea-ice coverage for the HAUSGARTEN area during the last 15 years; however, a time-series constructed from in situ measurements by Renner et al. (2014) exhibited a constant thinning of the

sea-ice by over 50% at the end of the melt season between 2003 and 2012.

Based on measurements by an array of oceanographic moorings at 78°50' N in the Fram Strait, Beszczynska-Möller et al. (2012) studied the variability in Atlantic Water (AW) temperature and volume transport in the West Spitsbergen Current (WSC). For the time period 1997–2010, they calculated a positive linear trend in the AW




Fig. 3. Water temperatures in the eastern Fram Strait between 1998 and 2014; annual means in the WSC at 250 m (data from Beszczynska-Möller et al., 2012; top) and at 2500 m water depth at the central HAUSGARTEN site off Svalbard (Glover et al., 2010 and Soltwedel, unpubl. data; bottom).

mean temperature of 0.06 °C per year, but no statistically significant trend was observed in the volume transport. Although comparing completely different water masses, annual mean temperatures at 250 and 2500 m water depth showed generally increasing values of about 1 °C at shallow depths and approx. 0.1 °C at the deep seafloor, respectively (Fig. 3).

Besides the overall trend, two Warm-Water Anomalies (WWA) were found in the AW passing through the Fram Strait between 1997 and 2010 (Beszczynska-Möller et al., 2012). The initial WWA occurred in 1999–2000, while the second, more pronounced one arose in late 2004 and continued until 2008. Whereas the first anomaly occurred in a confined area in the central Fram Strait, the latter one covered large areas in eastern parts of the strait, i.e., within the central HAUSGARTEN area. During the peak of the second warming (in September 2006), the anomalously warm AW, with temperatures >3 °C, even reached the East Greenland Current and the upper slope east of Greenland. On the eastern side of the WSC, the AW warmer than 3 °C was present at a depth of 250 m throughout the year, in contrast to earlier years when its occurrence was limited to the summer months (Beszczynska-Möller et al., 2012).

3. Time-series work at HAUSGARTEN

Time-series work in the open ocean and in the deep sea is logistically challenging due to the difficult accessibility of the area of interest, especially in polar regions. Thus, sampling is usually restricted and limited to small time windows when cruises by

Table 1

Key variables regularly assessed at the LTER observatory HAUSGARTEN.

Parameters	Pelagic zone	Sediment-water-interface	Sediments
Abiotic			
Temperature	×	×	
Oxygen	×	×	×
Currents (speed, direction)	×	×	
Sediment grain sizes, porosity			×
Biotic			
Phyto-/zooplankton (abundance, diversity)			
Particle flux rate (mainly phytodetritial matter)	×		
Biomarker in settling matter (marine, terrigenous)			×
Sediment-bound organic carbon (content, remineralization)		×	
Bacteria (activity, abundance, biomass, diversity)	×		
Meio-, macro-, megafauna (activity, abundance, biomass, diversity)			×
Demersal fish (activity, abundance, biomass, diversity)			×

research ships can be performed, finally resulting in snapshot-like datasets. Seasonally performed time-series from the open ocean and the deep sea are very rare. Such restriction can partly be overcome by using remote-sensing techniques, and by the installation of long-term (e.g., annually) moored systems carrying autonomous instruments allowing year-round measurements and sampling (see below). However, because of methodological restraints (e.g., restrictions to remote-sensing due to clouds, missing long-term stability for biochemical sensors, and preservation of samples in autonomous sampling devices) the application of these techniques still has its limitations.

One of the key characteristics of the scientific work at the LTER observatory HAUSGARTEN is the multidisciplinary approach, combining expertise from biologists, geochemists, sedimentologists, and physical oceanographers. International collaboration has been initiated in all fields where in-house experience was missing. An overview of key variables regularly assessed at HAUSGARTEN is presented in Table 1.

Water column studies at HAUSGARTEN include the assessment of physico-chemical parameters, plankton communities as well as flux measurements of particulate matter to the deep seafloor. An Autonomous Underwater Vehicle (AUV) equipped with water samplers and various sensors (e.g., a CTD as well as sensors for oxygen, nitrate, chlorophyll *a*, pCO₂, CDOM, and PAR) is used for upper water column studies (Wulff et al., 2013). Phyto- and zooplankton are sampled with water collectors and different nets (e.g., Multi-net, Bongo-net); analyses focus on species composition, abundance and spatial distribution. The settling of particles to the seafloor is assessed with conical sediment traps with an opening of 0.5 m² and equipped with 20 sampling cups (see Kremling et al., 1996). Samples are poisoned in situ with mercuric chloride for the preservation of samples during and after deployment. Treatment of the samples in the laboratory follows the description in Bodungen et al. (1991) and included the removal of so-called 'swimmers', i.e., zooplankton organisms that most probably enter the sediment traps actively (Harbison and Gilmer, 1986; Lee et al., 1989). Moorings and free-falling systems (bottom-lander), used as stationary platforms for observations and in situ experiments, are generally equipped with current meters and optodes to assess variations in hydrographic variables and oxygen concentrations, respectively.

Benthic communities are investigated over the entire size spectra from bacteria, meio- and macrofauna up to megafauna and demersal fish. Bacteria as well as meio- and macrobenthic organisms are sampled by multiple corer and box corer. Towed photo/video systems allow the non-invasive assessment of large-scale distribution patterns of epi/megabenthic organisms (for sampling procedures and sample processing, see Hoste et al. [2007], Budaeva et al. [2008], Bergmann et al. [2011], and Jacob et al. [2013]). Benthic studies at the HAUSGARTEN observatory also include biochemical analyses of sediment samples to estimate




Fig. 4. Monthly means of satellite-retrieved chlorophyll *a* from the GlobColour product (CHL) averaged for an area in the West Spitsbergen Current (76° N – 84° N and 0° – 15° E) for growth periods covering the months April through August between 1998 and 2012. The gray-shaded area indicates the period of the Warm-Water-Anomaly (WWA).

the input of phytodetrital matter to the seafloor as a food/energy source as well as to assess activities and biomasses of the small sediment-inhabiting biota (for analytical methods see Hasemann et al., 2013). Microprofiler and benthic chambers attached to free-falling systems (bottom-lander) are used to study remineralization rates and exchange processes at the sediment-water interface. Remotely Operated Vehicles (ROV) are routinely used at two-year intervals to carry out observations and targeted sampling, to position and service autonomous instruments, and to conduct in situ experiments at the deep seafloor.

The data presented in the following sections reflect the current state of progress in the processing of samples from the various compartments of the marine ecosystem at the LTER observatory HAUSGARTEN. For example, while results from the very labor-intensive meiofauna studies are only available until 2009, data sets from faster, straight-forward analyses cover also the most recent years. In any case, results presented below cover at least a decade of multidisciplinary time-series work at HAUSGARTEN.

3.1. Pelagic studies

In the upper pelagic system, the development of phytoplankton biomass was investigated via chlorophyll *a* concentrations derived from satellite data (April through August between 1998 and 2012) and chlorophyll *a* concentrations sampled and processed from discrete water depths in the euphotic zone during summer expeditions to the HAUSGARTEN observatory. The chlorophyll *a* concentrations obtained with remote-sensing observations by the NASA sensors SeaWiFS and MODIS, and the ESA sensor MERIS, were merged to one CHL data product (GlobColour product, see <http://hermes.acri.fr>). Beside seasonal variability, chlorophyll *a* in the West Spitsbergen Current (WSC) during the years 1998–2007 stayed relatively constant; however, an overall increase of the chlorophyll *a* biomass between 2008 and 2012 is visible in the satellite data (Fig. 4) as well as in the water column (data not shown), pointing to a phytoplankton biomass increase following the WWA in the Fram Strait between 2005 and 2008.

Cherkasheva et al. (2014) found no direct link between sea-ice concentrations and chlorophyll *a* concentrations for the wider Fram Strait region between 1998 and 2009. Therefore, the authors considered additional physical factors potentially affecting phytoplankton distribution, including sea-ice thickness, water temperatures, and salinity gradients from a validated sea-ice ocean circulation model (NAOSIM); the latter data were then used to calculate the time-series of stratification within this region. Results

showed that the Marginal Ice Zone (MIZ) was promoting phytoplankton growth by stratifying the water column. For the years 1998 through 2009, highest mean chlorophyll concentrations in the MIZ reached up to $0.8 \text{ mg CHL m}^{-3}$ (average values for the productive season April–August). In the open ocean, the phytoplankton growth correlated best with thermal stratification formed by solar heating of the upper ocean layers (Cherkasheva et al., 2014). The coastal zone around Svalbard showed processes associated with the presence of landfast ice that is suppressing rather than promoting the phytoplankton growth. During the 12 years of observations (1998–2009), chlorophyll concentrations increased in the southern part of the Fram Strait, associated with an increase in sea-surface temperature and a decrease in Svalbard coastal ice (Cherkasheva et al., 2014).

The species composition of unicellular plankton organisms has been determined by collecting water samples from the chlorophyll *a* maximum layer at the HAUSGARTEN central station during summer months between 1998 and 2011 (Mebrahtom Kidane, 2011). In the years 1998 and 2003, diatoms of the genus *Thalassiosira*, *Chaetoceros*, and *Fragilariaopsis* dominated the phytoplankton (Fig. 5).

These years were characterized by longer periods of sea-ice coverage and comparably cold temperatures in surface waters. Coccolithophores, mainly composed of *Emiliania huxleyi*, prevailed in the samples from 2004. Throughout the following years, i.e., during and after the WWA, phytoplankton was dominated by the prymnesiophyte *Phaeocystis pouchetii*. Concomitant to the shift in phytoplankton composition, we noted an increase in protozooplankton abundance (e.g., ciliates and tintinnids) in the water samples. The observed changes in the species composition of unicellular planktonic protists is also partly reflected in the composition of the settling matter in sediment traps at the central HAUSGARTEN site, indicating modifications also in element cycling in the Arctic Ocean (see Section 3.2).

3.2. Flux studies

Results from sediment trap deployments indicate a co-variation of the total particulate matter flux and the share of biogenic matter within the overall flux with the prevailing ice conditions. Fluxes of total matter and components thereof (i.e., carbonate, particulate organic carbon and nitrogen, biogenic particulate silica and biomarkers, e.g., brassicasterol, campesterol, β -sitosterol, long-chain alkenones, IP₂₅) revealed a bimodal seasonal pattern showing elevated sedimentation rates during May/June and at the end of




Fig. 5. Composition of unicellular planktonic protists ($>3 \mu\text{m}$) in the chlorophyll *a* maximum of the water column at the central HAUSGARTEN site for eight years from 1998 to 2011.

the growth season in August/September (Bauerfeind et al., 2009; Lalande et al., 2013).

The total dry weight (DW) of settling matter over the years was rather constant, except for 2003 as well as 2008 and 2009, where DW values were slightly elevated (data not shown). The export flux of particulate organic carbon (POC) was rather low ($2\text{--}3 \text{ g m}^{-2} \text{ y}^{-1}$) suggesting a large retention and recycling of the POC in the water above 200–300 m. Unexpectedly, fluxes of POC in the upper water column did not change significantly between 2000 and 2012, whereas the flux of biogenic particulate silica, a proxy for diatoms, decreased markedly from ~ 1.5 to $<0.5 \text{ g m}^{-2} \text{ y}^{-1}$, thereby indicating a shift in the dominant primary producers toward non-diatom organisms (Bauerfeind et al., 2009; Lalande et al., 2013). The annual flux of all biogenic components (i.e., opal+organic carbon+carbonate) between 2000 and 2012 varied between 8.6 and $15.7 \text{ g m}^{-2} \text{ y}^{-1}$, and was almost constant before 2008/09, but increased following the WWA (Fig. 6, top).

The relative contribution of the biogenic components indicate that the proportion of opal decreased markedly to 10% or less, whereas the proportion of carbonate increased from 20% to 30% in the first years to >50% after the WWA (Fig. 6, bottom). The

proportion of organic carbon varied from 30% to 60% and generally showed no clear trend between 2000 and 2012. An exception was the period 2009/2010 when carbonate and organic carbon contributed with $\sim 30\%$ and $\sim 60\%$, respectively, to the flux of biogenic matter. However, this may be explained by the deviating depth of the sediment trap which, by accident, was installed ~ 120 m shallower compared to the other years.

The annual fluxes of brassicasterol, a biomarker for phytoplankton of marine and/or fluvial origin (Fahl and Nöthig, 2007; Fahl and Stein, 2012 and references therein), showed constant values, except for 2006/2007, where brassicasterol concentrations in the settling matter were clearly higher (Fig. 7, top).

The overall persistent flux over the last years may indicate a mixed signal related to phytoplankton input from the water column and ice-transported material via the Transpolar Drift. In contrast, biomarkers for terrigenous origin (i.e., campesterol and β -sitosterol) decreased continuously between 2000 and 2009 (Fig. 7, bottom). A similar trend was found for the biomarker IP₂₅ (data not shown), a highly branched isoprenoid with 25 carbons, which is biosynthesized exclusively by diatoms living in arctic sea-ice (Belt et al., 2007; Müller et al., 2009). Considering that the




Fig. 6. Annual flux of biogenic matter (top) and percentage composition of its components (bottom) between 2000 and 2012 (n.d. = no data). The gray-shaded area indicates the period of the WWA.




Fig. 7. Annual flux of the biomarker brassicasterol (top) and the sum of campesterol and β -sitosterol (bottom) for seven years between 2000 and 2009. The gray-shaded area indicates the period of the WWA.

terrigenous organic matter in the HAUSGARTEN area is derived by ice transport from the arctic marginal seas (e.g., the Laptev Sea), the good correlation between the decreasing fluxes of IP₂₅ (a sea-ice marker) and the terrigenous sterols campesterol and β -sitosterol is probably indicative of a generally decreasing sea-ice cover and thinner ice in the central Arctic Ocean.

Differences in the export of protist assemblages during the period of highest flux in September were assessed by using the Terminal Restriction Fragment Length Polymorphism (T-RFLP) method (Marsh, 1999), which is based on sequence heterogeneity, e.g., within the 18S rDNA of different taxa. The composition of differently-sized restriction fragments in a sample acts as a characteristic fingerprint of a microbial community that allows qualitative comparisons of community compositions. A multidimensional scaling (metaMDS) plot based on T-RFLP differences in the exported protist assemblages at the central HAUSGARTEN site (79° N, 04° E) between 2000 and 2010 suggests that the community profiles segregate into four groups (Fig. 8). Group one is composed of samples collected in 2000–2002, group two in 2004–2005, group three in 2006–2007, and group four in 2008–2010. An ANOSIM analysis to test the significance of the a priori defined groups resulted in an *R*-value of 0.9256 and a *p*-value of 0.001, indicating highly significant differences between the defined groups. The metaMDS plot also implies that the protist communities in 2006–2010 were more similar to each other than to those of the years before the WWA (Fig. 8).

Valuable information about the composition of the zooplankton community in the HAUSGARTEN area can be achieved by analyzing the composition of swimmers that are caught as a bycatch in the sediment traps (e.g., Seiler and Brandt, 1997; Willis et al., 2008; Kraft et al., 2011; Bauerfeind et al., 2014). At the HAUSGARTEN, we analyzed changes in the abundance of two major groups, i.e., pelagic amphipods (species of the genus *Themisto*) and gastropods (thecosome pteropods of the genus *Limacina*).

Within the amphipods, three *Themisto* species were observed regularly and in great abundance, i.e., the arctic species *Themisto libellula*, the sub-arctic *T. abyssorum*, and the North Atlantic species *T. compressa* (Kraft et al., 2012). The pteropod species consisted of the polar species *Limacina helicina* and the transitional,




Fig. 8. Multidimensional scaling (metaMDS) plot to illustrate differences between the 18S T-RFLP patterns of the different protist communities in the sediment traps. The plot is based on the calculation of Jaccard indices for the data set derived from amplification of an 18S rDNA fragment and subsequent digestion with the restriction enzyme HaeIII.

boreal warm water-adapted *Limacina retroversa* (Bauerfeind et al., 2014). The occurrence and composition of these organisms in the samples underwent substantial changes during the period 2000–2012 (Fig. 9). Pteropod numbers increased significantly during the WWA, showed comparably low values in 2009/2010, but steeply increased again until the end of the time-series (Bauerfeind et al., 2014; Busch et al., 2015). The overall rise in pteropods numbers was accompanied by a switch in dominance of species from the cold-adapted *L. helicina* that dominated in the first years until 2004/2005 toward the warm water-species *L. retroversa* in the subsequent years (Fig. 9, top).

The number of amphipods also increased steadily (Fig. 9, bottom), mainly due to increased numbers of the sub-arctic species *T. abyssorum*. The arctic species *T. libellula*, on the other hand, did not follow the general trend after the WWA, but was present in varying abundances during all the years since 2000. Most remarkably, we also noted the occurrence of *T. compressa*, an amphipod species, which has not been recorded in the Fram Strait before. This species was first observed sporadically during 2001–2005, and was present continuously in the subsequent years with largest numbers in 2011, when also juveniles of the species were recorded for the first time (Kraft et al., 2013).

3.3. Benthic studies

3.3.1. Biogenic sediment compounds

Benthic long-term studies at HAUSGARTEN include the analyses of biogenic sediment compounds to estimate food availability at the deep seafloor and the biomass of the small benthic biota. The input of phytodetritus, representing the main food source for benthic organisms, is estimated by analyzing concentrations of sediment-bound chloroplastic pigments (Yentsch and Menzel, 1963; Holm-Hansen et al., 1965). As the fluorometric method used in our studies does not discriminate between individual breakdown products, we prefer the term chloroplastic pigment equivalents (CPE) for the bulk pigments analyzed (Thiel, 1978). Benthic biomass within the sediments was determined by different biochemical assays. Particulate proteins, indicating the bulk of “living” and “dead biomass”, i.e., small organisms and detrital matter,




Fig. 9. Abundance of *Limacina* pteropods (top) and *Themisto* amphipods (bottom) in the sediment traps at the central HAUSGARTEN site (n.d. = no data). The gray-shaded area indicates the period of the WWA.

respectively, were analyzed photometrically (Greiser and Faubel, 1988). The analysis of sediment-bound phospholipids (indicative of cellular membranes) provides biomass estimates for benthic micro-organisms (Findlay et al., 1989). According to the sample sizes used for these studies, biomass estimations cover the entire small-sized benthic biota, including bacteria, fungi, protozoans, and the metazoan meiofauna.

Spatial and temporal assessments of biogenic sediment compounds along the bathymetric HAUSGARTEN transect off Svalbard showed generally decreasing values with increasing water depth, but also distinct interannual variations between 2000 and 2010. Hovmöller diagrams displaying anomalies from ten year means reveal a sharp increase in sediment-bound pigment concentration, i.e., phytodetrital matter between 2006 and 2007 (Fig. 10, left).

At the same time, during the peak of the WWA, a similar but less pronounced shift to positive values was found for particulate proteins (total biomass), whereas phospholipid concentrations (microbial biomass) followed the trend only vaguely. While strong positive values were found at the onset of the time-series study (where pigment and protein concentrations were already low), enhanced microbial biomass values since 2007 were only found for certain water depths (Fig. 10, right). Differences in food quality at the beginning of the time-series and after the increased phytodetrital matter availability at the deep seafloor following the WWA might obscure a clear relationship.

3.3.2. Life at the deep seafloor

Benthic bacteria at HAUSGARTEN were assessed using a LIVE/DEAD® BacLight™ Bacterial Viability Kit protocol adapted to marine sediments (Quéric et al., 2004). Along the bathymetric transect off Svalbard, the analyses revealed a decrease of both bacterial viability and abundance with increasing water depth. Maximum bacterial abundance ($4.9\text{--}5.4 \times 10^8$ cells cm^{-3}) was found in the uppermost sediment layer at ~ 1250 m water depth; lowest bacterial cell numbers ($0.8\text{--}1.2 \times 10^8$ cells cm^{-3}) were observed at the deepest stations (~ 5600 m water depth). The percentage of viable (and thus potentially active) cells ranged between 20% and 60% within the first, and 10–40% within the fifth centimeter of sediment throughout the transect (Quéric et al., 2004).

Regional distribution patterns of the benthic bacterial community at HAUSGARTEN were examined by Jacob et al. (2013). Samples from 13 stations were retrieved from the bathymetric (1284–3535 m water depth, 54 km in length) and the latitudinal transect (~ 2500 m water depth; ~ 120 km in length) (see Fig. 1). Applying the 454 Massively Parallel Tag Sequencing (MPTS) method, Jacob et al. (2013) found a spatially highly diverse bacterial community, with strong water depth-related patterns in community structure along the bathymetric transect, whereas no significant changes in community structure were observed with increasing spatial distance along the latitudinal transect. MPTS identified a total of 41 phyla, 78 classes, 136 orders, 215 families, and 410 genera. Most of the bacteria belonged




Fig. 10. Spatio-temporal diagrams of interannual variability in different biogenic sediment compounds along the bathymetric transect off Svalbard between 2000 and 2010: anomalies from the long-term mean in sediment-bound chloroplastic pigments (CPE), particulate proteins (PROT), and phospholipids (LIPIDS); blue colors = lower values, white and red colors = values exceeding the long-term mean.

to the phylum Proteobacteria (47%), with the most abundant classes being Gammaproteobacteria (23%), Deltaproteobacteria (15%), and Alphaproteobacteria (7%). The second most abundant phylum was Bacteroidetes (9%), including the classes Flavobacteria (3%) and Sphingobacteria (5%). Other abundant phyla were Actinobacteria (3%), Acidobacteria (5%), and Verrucomicrobia (4%). The richness of operational taxonomic units decreased substantially during the WWA in 2005–2007, and again increased in 2008 to a similar level as in 2003 (Jacob, 2014). Changes in the bacterial community were attributed to changes in food availability along the bathymetric transect as well as during the WWA (see Section 3.3).

Meiobenthic studies at HAUSGARTEN between 2000 and 2009 focused on the metazoan community and revealed densities ranging from 269 ± 119 to 3248 ± 22 ind. 10 cm^{-2} (Hoste et al., 2007; Grzelak, 2015). Nematodes dominated the metazoan meiofauna at every depth and sampling year (85–99%), followed by harpacticoid copepods (up to 4.6%). The expected pattern of gradually decreasing meiobenthic densities with increasing water depth was not documented. Instead, the bathymetric transect at HAUSGARTEN could be roughly subdivided into a shallow part with both high nematode and copepod densities between approximately 1000 and 2000 m water depth (means \pm SD: 2259 ± 157 nematodes 10 cm^{-2} , and 50 ± 4 copepods 10 cm^{-2}), and a deeper part between approximately 3000 and 5500 m water depth with clearly lower nematode and copepod densities (means \pm SD: 595 ± 52 nematodes 10 cm^{-2} , and 11 ± 2 copepods 10 cm^{-2}). Compared to deep-sea studies from the North Atlantic, the nematode community at HAUSGARTEN is characterized by very high species numbers, even though the number of genera was in the same order of magnitude (Hoste, 2007). Rare genera and species were diverse, but represented <1% of the total nematode abundance. The overall nematode community structure at HAUSGARTEN was similar to other typical

deep-sea communities, with dominant genera such as *Tricoma*, *Desmoscolex*, *Acantholaimus*, *Halalaimus* and members of the Monhysteridae. Except for the deepest stations, deposit feeders were dominant at each water depth followed by epistrate feeders. The nematode community composition changed gradually with water depth. Differences were mainly due to variations in the relative abundance of genera rather than the presence/absence of genera. At the species level, some indicative species could be found at most water depths, especially within the Monhysteridae, Xyalidae, and Desmoscolecidae. Aside from water depth as the main factor accounting for differences in nematode communities, interannual variability in nematode community structure was apparent with high relative abundances of *Dichromadora*, *Microlaimus*, *Acantholaimus*, *Halalaimus*, *Daptoneema*, and *Tricoma* related to enhanced food availability (Hoste, 2007). Since 2005, *Sabatieria* and *Sphaerolaimus* replaced *Microlaimus* and *Daptoneema* in the list of the most abundant genera (Grzelak, 2015).

Combined data from Hoste et al. (2007) and Grzelak (2015) for the central HAUSGARTEN site between 2000 and 2009 showed an overall decrease in nematode densities from 2002 to 2005 and generally increased values in the subsequent years, with highest values in 2006 (Fig. 11). Moreover, based on the classification of the nematode communities into feeding types, we observed overall increasing trophic diversities (specified by the Index of Trophic Diversity, ITD, where lower values indicate higher diversity) at the central HAUSGARTEN station between 2000 and 2006, with highest diversity in 2006. Most probably affected by changes in food availability during the WWA (see Section 3.3), trophic diversity of the nematode communities clearly decreased in 2007, with an increasing trend in the following two years. The comparably high trophic diversity of the nematode communities in 2006 can be traced back to almost similar proportions of particle and epigrowth feeders together with a comparatively high proportion of




Fig. 11. Mean nematode densities and standard deviations from three replicates (bars), and the index of trophic diversity (bullets) of the nematode community inhabiting sediments from the central HAUSGARTEN site (~ 2500 m water depth) off Svalbard from 2000 to 2009.

predators/scavengers and omnivore feeding nematodes. The low trophic diversity in 2007 is mainly caused by the high dominance of non-selective particle feeders (up to 48%), whereas similar proportions of selective and non-selective particle feeders (between 31% and 37%) as well as comparatively high proportions of predators/scavengers and omnivore feeding nematodes (between 8% and 11%) lead to the increasing trophic diversity in the years 2008 and 2009.

Macrofauna were only irregularly sampled at HAUSGARTEN, so far not allowing the study of temporal variations in the area. However, spatial studies revealed distinct depth-related, but also horizontal differences in macrofaunal community structure.

Along a transect toward the Kongsfjorden (Svalbard), Włodarska-Kowalcuk et al. (2004) found a community replacement along the depth gradient with four different types of assemblages categorized as SHELF (<370 m), SHALLOW SLOPE (~ 500 m), DEEP SLOPE (~ 1500 m), and RISE (>2000 m). The dominant macrofauna taxa in different depth zones along the bathymetric transect are summarized in Table 2.

Macrofaunal species richness (expressed by the number of species per sample) decreased with water depth. There was, however, no clear bathymetric pattern in diversity measures; the classic increase of species richness and diversity at mid-slope depths was not observed. According to Włodarska-Kowalcuk et al. (2004), Shannon-Wiener diversity was the highest at 525 m water depth ($H'_{(\log e)} = 3.54$) and lowest at 2025 m depth ($H'_{(\log e)} = 1.46$). Species richness and diversity at the deep stations were much lower than in comparable studies from the temperate North Atlantic. Włodarska-Kowalcuk et al. (2004) related this finding to the geographic

isolation of Greenland–Icelandic–Norwegian Seas from the Atlantic pool of species. Macrofaunal biomass was negatively correlated with depth ($R = -0.86$), and ranged from 61 g ww m^{-2} at the shallowest station sampled (212 m) to 1 g ww m^{-2} at 2025 m water depth. Biomasses were much higher than at similar depths in the multi-year ice-covered High Arctic (Kröncke, 1994, 1998), while resembling those from temperate and tropical localities (Flach et al., 1999; Galeron et al., 2000).

In their study on horizontal community patterns, Budaeva et al. (2008) used nested sampling to study different scales of macrofauna distribution at three sampling sites along a 26-km long transect at 2500 m close to the central HAUSGARTEN site. Two separate macrofaunal assemblages were identified. The first assemblage comprised individuals found in three box core samples taken at one of the three sites. The second assemblage contained all individuals collected with the other six box corers taken at the remaining two sites located about 2 km away from each other and 23 km away from the third site. Budaeva et al. (2008) compared their findings with the results published by Włodarska-Kowalcuk et al. (2004) and concluded that the macrobenthos in the HAUSGARTEN region appears to be organized at three spatial scales: communities that replace each other along the depth gradients, species assemblages that contribute to the heterogeneity within each vertical zone, and single-species patches that create variability at the local scale (Budaeva et al., 2008).

Large-scale distribution patterns of epi/megafauna organisms are regularly assessed by a towed photo/video system (Ocean Floor Observation System, OFOS). Investigations along the depth transect off Svalbard revealed a clear bathymetric zonation of megafaunal assemblages (Soltwedel et al., 2009). Megafaunal densities ranged between 11 and 38 ind. m^{-2} with highest densities at 1650 m and lowest densities at 3000 m depth. The number of taxa ranged between 4 at 5500 m, and 27 at 1650 m water depth. Deposit-feeding brittle stars (*Ophiocent cf. hastatum*) dominated the continental margins up to depths of ~ 1300 m with mean densities of 17 ind. m^{-2} . At a station only 300 m deeper, small yet unidentified amphipods dominated the fauna (37 ind. m^{-2}). At ~ 2500 m water depth, deposit-feeding sea cucumbers (*Kolga hyalina*) and suspension-feeding sea anemones accounted for the majority of megafauna (4 and 3 ind. m^{-2} , respectively). Suspension-feeding crinoids (*Bathycriinus carpenterii*, 5 ind. m^{-2}) and sea anemones (2 ind. m^{-2}) were the most abundant groups recorded at ~ 3000 m water depth. Small arctic sea cucumbers (*Elpidia heckeri*, 30 ind. m^{-2}) dominated the Molloy Hole. The zonation patterns appear to be controlled primarily by food availability (Soltwedel et al., 2009). Log₂-based Shannon-Wiener diversity ranged between 0.34 and 2.58 with highest densities at about 2620 m ($H'_{(\log 2)} = 2.58$) and between 2965 and 3043 m ($H'_{(\log 2)} = 2.50$) water depths. Diversity was lowest at 1642 to 1688 m ($H'_{(\log 2)} = 0.34$) and between 5333 and 5404 m ($H'_{(\log 2)} = 0.51$).

Table 2

Dominant macrofauna taxa and proportion from the entire macrofauna community (%) at different depth zones along the bathymetric transect in the eastern Fram Strait; P = Polychaeta, C = Crustacea, B = Bivalvia, E = Echinodermata, S = Sipunculida.

Shelf (<370 m)	Shallow slope (~500 m)			Deep slope (~1500 m)			Rise (>2000 m)		
<i>Prionospio cirrifera</i>	23.5%	P	<i>Yoldiella lucida</i>	8.9%	B	<i>Thyasira dubnari</i>	22.5%	B	<i>Myriochele heeri</i>
<i>Cossura longocirrata</i>	17.7%	P	<i>Exogone cf. verugera</i>	8.5%	P	<i>Aricidea</i> spp.	16.8%	P	<i>Myriochele fragilis</i>
<i>Lumbrineris</i> sp.	12.5%	P	<i>Leucon nathorsti</i>	5.1%	C	<i>Paraonis/Paradoneis</i> sp.	5.6%	P	<i>Paraonis/Paradoneis</i> sp.
<i>Leitoscoloplos</i> sp.	10.2%	P	<i>Chone paucibranchiata</i>	4.7%	P	<i>Ophiura</i> juv.	4.6%	E	<i>Notomastus</i> sp.
Chaetozone group	7.7%	P	<i>Tanaidaceae</i> sp. A	4.3%	C	<i>Myriochele fragilis</i>	3.5%	P	<i>Sipunculida</i> nd
<i>Levinsenia gracialis</i>	3.5%	P	<i>Nicomache</i> sp.	4.1%	P	<i>Ilyarachnidae</i> nd	3.2%	C	<i>Tanaidacea</i> sp. A
<i>Ophiura robusta</i>	3.0%	E	<i>Tanaidacea</i> sp. B	3.9%	C	<i>Sipunculida</i> nd	3.2%	S	<i>Haustoriidae</i> nd
<i>Heteromastus filiformis</i>	2.9%	P	<i>Nothria conchylegaa</i>	3.7%	P	<i>Capitellidae</i> nd	2.8%	P	<i>Yoldiella</i> sp.
<i>Myriochele oculata</i>	1.5%	P	<i>Heteromastus filiformis</i>	3.0%	P	<i>Oedicerotidae</i> nd	2.5%	C	<i>Aricidea</i> spp.
<i>Maldane sarsi</i>	1.0%	P	<i>Ophiura</i> juv.	3.0%	E	<i>Tanaidacea</i> sp. A	1.8%	C	<i>Ilyarachnidae</i> nd




Fig. 12. Mean megafaunal densities in different feeding groups at ~1250 m (top) and ~2500 m water depth (bottom) between 2002 and 2012, and 2011, respectively. (n.d.= no data).

Analysis of seafloor photographs taken at ~1250 and ~2500 m water depth off Svalbard in the years 2002, 2004, 2007, and 2011/2012 revealed significant temporal variation in megafaunal densities and community composition (Fig. 12).

At the shallower site, megafaunal density was similar in 2002 and 2007, but significantly higher in 2012 (Kruskall-Wallis test, pairwise Mann-Whitney tests applying a Bonferroni correction of $p=0.017$; Fig. 12, top) (Meyer et al., 2013). OFOS footage taken at 2500 m water depth indicated a significant gradual decrease in total megafaunal densities between 2002 and 2007 (Bergmann et al., 2011). While suspension-feeder densities also declined, predator/scavenger and deposit-feeder densities decreased to such a degree that suspension feeders made up >90% of the megafauna in 2007. After 2007, however, total megafaunal densities grew again verging on initial levels in 2011 (Fig. 12, bottom) (see also Müller et al., this issue). Overall, megafaunal densities from 2002 and 2011 were significantly higher than those recorded in 2004 and 2007 (Kruskall-Wallis test, pairwise Mann-Whitney tests with a Bonferroni correction $p<0.001$).

4. Insights from 15 years of observations at HAUSGARTEN

HAUSGARTEN is synonymous with the longest multidisciplinary time-series study at a polar open-ocean site. Multidisciplinary studies at the LTER observatory covered large parts of the marine ecosystem and already exhibited a number of interesting temporal trends.

A warming event in the Atlantic Water layer in the eastern Fram Strait between 2005 and 2008 left a strong “footprint” in the arctic marine ecosystem in the HAUSGARTEN area. Impacts of the Warm-Water Anomaly (WWA) encompassed the increased growth of sub-arctic primary producers as well as a northward displacement, increased abundance and reproductive output of sub-arctic species, leading finally to changes in community structure due to

range shifts. Cascading effects propagating through the entire food web led to an astoundingly rapid and far-reaching reaction of the entire open-ocean ecosystem from the pelagic to the deep seafloor. A number of variables monitored at the HAUSGARTEN observatory seem to fall back to their previous state subsequent to the WWA (e.g., decreasing proportions of *P. pouchetii* between 2009 and 2011), while others seem to remain at a different level (e.g., constantly low amounts of diatoms and high densities of the sub-polar pteropod species *L. retroversa* as well as the enduring occurrence of the North Atlantic amphipod species *T. compressa*). At least certain components of the marine ecosystem in the Fram Strait might have reached a “tipping point”, i.e., a threshold in environmental forcing at which the future state of the system is qualitatively altered (Lenton et al., 2008).

It is a common knowledge that the Marginal Ice Zones (MIZ) are biologically very important (Sakshaug and Skjoldal, 1989), because stratification due to ice melting combined with a generally ample supply of nutrients sets the stage for vigorous phytoplankton production in the upper water layers. Zooplankton, pelagic fish, marine mammals and seabirds aggregate at the ice edge and exploit phytoplankton production in the MIZ. Large parts of the biological production in the MIZ sinks through the water column and forms a basis for rich communities of benthic organisms. Data collected at HAUSGARTEN showed that the proximity of the sea-ice in northern and western parts of the Fram Strait positively affects the flux of particulate matter to the seafloor. In 2003 as well as in 2008 and 2009, when ice anomalies showed strong positive values, material collected in sediment traps at the central HAUSGARTEN site showed increased total dry weights which, in this case, could be attributed to increased amounts of lithogenic matter settling out during ice melt (Bauerfeind et al., 2009). Overall increased flux rates in biogenic particulate silica (a proxy for diatoms) and zooplankton fecal pellets in the presence of the ice cover at HAUSGARTEN were detected by Lalande et al. (2013). A close correlation between maximum sedimentation rates and the position of the ice edge was also shown by Ramseier et al. (1999) for mooring sites in the north-eastern Greenland Sea.

Although sea-ice has the inherent potential to influence physico-chemical and biological processes in the water column (see above), most pronounced variations in the marine ecosystem at HAUSGARTEN appeared to be related to large-scale temporal variations in surface water temperature. The warm water pulse (WWA) detected between 2005 and 2008 (Beszczynska-Möller et al., 2012) led to significant changes in planktonic protist assemblages and in unicellular phytoplankton communities (Mebratoom Kidane, 2011). While macroscopic and biochemical analyses on sediment trap material revealed an overall decrease in diatoms since the beginning of the time-series studies at HAUSGARTEN, the WWA induced an abrupt shift in phytoplankton to a community dominated by micro- and nanoflagellates. A general shift from a dominance of diatoms to a dominance of picoplankton in a warmer Arctic Ocean is commonly predicted (Morán et al., 2009) and already supported by observations in the central Arctic Ocean (Li et al., 2009; Zhang et al., 2015).

Increasing concentrations of carbonate in the sediment trap material at the central HAUSGARTEN site during and after the WWA are due to the strong increase in pteropod densities (Bauerfeind et al., 2014). The sudden mass occurrence of the sub-polar pteropod species *L. retroversa* reflects a northward displacement (range shift) of sub-arctic and temperate species. Still enhanced abundances of *L. retroversa* in years following the WWA (Busch et al., 2015) might point to an increased reproductive output of this originally sub-arctic species further south and a subsequent transport to the Fram Strait with the prevailing currents. A northward shift in the latitudinal distribution of boreal species was also found for other mesozooplankton organisms in the epipelagic zone of the

West Spitsbergen Current (Weydmann et al., 2014). Between 2001 and 2009, Weydmann et al. (2014) observed a smooth change in dominance of ubiquitous and boreal/arctic taxa such as *Oithona similis* and *Triconia* sp. toward a stronger dominance of boreal taxa such as *Calanus finmarchicus*.

A similar range shift was found for pelagic amphipods, whereby the sub-arctic species *Themisto abyssorum* and the North Atlantic species *T. compressa* increasingly replaced the arctic species *T. libellula* during the WWA (Kraft et al., 2012). In fact, for Kongsfjorden (Svalbard), Hop et al. (2006) noticed the replacement of *T. libellula* by *T. abyssorum* already for the time period 1996–2002. Indicated by the presence of a complete temporal series of developmental stages, Kraft et al. (2013) found first evidence of reproductive success in the boreal amphipod species *T. compressa* at HAUSGARTEN observatory in 2011. Warm-water pulses in the WSC, like the WWA observed between 2005 and 2008, might also be responsible for modern incursions of tropical Radiolaria into the Arctic Ocean, as detected by Bjørklund et al. (2012) in areas north of Svalbard.

Water temperature-related changes in the composition of the phyto- and zooplankton communities in the HAUSGARTEN area alter the flux of particulate matter to the seafloor with accompanying impacts on the entire food web and the marine biogeochemistry (Wohlers et al., 2009). Significantly increased concentrations in biogenic sediment components indicating the input of organic matter to the seafloor and the biomass of the small benthic biota during the WWA confirmed changes in the quantity and quality of the settling matter. Apparently, these changes directly impacted the entire benthic communities from bacteria to megafauna.

Changes in benthic bacterial communities along the bathymetric HAUSGARTEN transect off Svalbard were mostly driven by changes in the overall food availability (Jacob et al., 2013), whereas food quality is likely to affect the meiobenthic nematode communities. Although the trophic structure of the nematode communities is simple, and nematodes probably compete for the same bacterial-based food source, a similar amount of food of different quality may well have different effects on the community and could explain differences. According to Soetaert and Heip (1995), increased and high quality food supply could allow for a higher trophic complexity to develop within nematode communities. When the supply of organic matter to the bottom is high, a size-based food chain can develop within the nematode communities and predatory nematodes can become an important part of the food web (Soetaert and Heip, 1995). This might explain the comparably high proportions of predators/scavengers and omnivorous nematodes at the central HAUSGARTEN site in the years following the increased food availability in 2006/2007.

While time-series studies on bacterial and meiofauna (nematode) assemblages at the HAUSGARTEN observatory showed an immediate response to changes in food availability at the deep seafloor, megafaunal communities apparently reacted with some delay. In fact, lowest megafaunal densities were found in 2007, when biogenic sediment compounds already indicated increased food availability at the deep seafloor. However, a 1–1.5-yr time lag in the response of the deep-sea megafauna to food availability, as observed by Ruhl (2007) for megafaunal assemblages in the deep North-East Pacific, could explain the significantly higher megafaunal densities in the eastern Fram Strait in 2011 and 2012. Interestingly, variations in megafaunal densities at HAUSGARTEN also coincided with significant changes in the trophic structure of the megafaunal community.

The major forcing on today's arctic environment is anthropogenic climate change, which is responsible for the recent increase in atmospheric and ocean temperatures over the Arctic (IPCC, 2013). However, even after 15 years of intense

time-series work at HAUSGARTEN, we cannot yet judge with complete certainty whether the observed changes in environmental parameters and the marine biota indicate lasting alterations due to anthropologically-induced global changes, or whether they reflect natural variability on multiyear time-scales, for example, in relation to decadal oscillatory atmospheric processes expressed by indices such as the Atlantic Multidecadal Overturning (AMO), the North Atlantic Oscillation (NAO), and the Arctic Oscillation (AO) (e.g., Drinkwater et al., 2014).

Although we cannot conclude with absolute confidence that the WWA occurring in the Fram Strait between 2005 and 2008 is part of a broad human-induced shift toward a warmer Arctic Ocean, the observed alterations due to this warm water pulse may be seen as a "blueprint" for how the marine ecosystem might develop in an overall warming Arctic Ocean. However, apart from atmospheric and oceanic warming, the Arctic is also characterized by the presence of a number of other tipping elements (Duarte et al., 2012), including decreasing sea-ice coverage modifying heat fluxes and gas exchanges (Serreze et al., 2007), increased freshwater discharge from melting permafrost (Peterson et al., 2002), ocean acidification reducing CO₂ storage and calcification in organisms (Steinacher et al., 2009), and human activities increasing the risk of pollution (Huntington et al., 2007; Bergmann and Klages, 2012; Obbard et al., 2014).

Long time-series studies remain the key tools to provide insights into processes and dynamics within an arctic marine ecosystem, and in fact, there is an urgent need for heightening efforts to study marine ecological variability and changes induced by climate change in the Arctic (Wassmann et al., 2011). Only long-term studies on seasonal to multi-decadal time scales will allow for improved predictions of the temporal development of the marine ecosystem in a future Arctic Ocean.

Acknowledgements

We wish to thank the officers and crews of RVs *Polarstern*, *Maria S. Merian*, and *L'Atalante* for their help during expeditions to the LTER observatory HAUSGARTEN. We gratefully acknowledge the tremendous support of our engineers, technicians and student workers in the laboratory and at sea over all the years. The remote sensing part of this study was supported by POLMAR Helmholtz Graduate School for Polar and Marine Research and Helmholtz Impulse and Network Fond at the Alfred Wegener Institute (project PHYTOOPTICS). Satellite data was provided by NASA (SeaWiFS, MODIS) and ESA (MERIS, GlobColour). Financial support was given to MJ through the ERC Advanced Grant ABYSS (no. 294757). MWK was financially supported by the Polish National Science Centre project funded based on decision no. DEC-2012/05/B/NZ8/02654. KG was supported by the Polish National Science Centre grant 1721/B/P01/2011/40. CH was supported by the BFMB-Project TRANSDRIFT ("System Laptevsee" – Das transpolare System des Nordpolarmeeres). MB was funded by the Helmholtz Alliance ROBEX (Robotic Exploration of Extreme Environments). KM's work was supported by a Fulbright Study-Research Grant. This is publication "e37840" of the Alfred Wegener Institute, Helmholtz-Centre for Polar and Marine Research, Bremerhaven, Germany.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ecolind.2015.10.001>. These data include Google maps of the most important areas as described in this article.

References

- AMAP, 2013. AMAP Assessment 2013: Arctic Ocean Acidification. Arctic Monitoring and Assessment Programme, Oslo, Norway, viii + 99 pp.
- Arrigo, K.R., van Dijken, G., Pabi, S., 2008. Impact of a shrinking Arctic ice cover on marine primary production. *Geophys. Res. Lett.* 35 (19), L19603.
- Arrigo, K.R., Perovich, D.K., Pickart, R.S., Brown, Z.W., van Dijken, G.L., Lowry, K.E., Mills, M.M., Palmer, M.A., Balch, W.M., Bahr, F., Bates, N.R., Benitez-Nelson, C., Bowler, B., Brownlee, E., Ehn, J.K., Frey, K.E., Garley, R., Laney, S.R., Lubelczyk, L., Mathis, J., Matsuoaka, A., Mitchell, B.G., Moore, G.W.K., Ortega-Retuerta, E., Pal, S., Polashenski, C., Reynolds, R.A., Schieber, B., Sosik, H.M., Stephens, M., Swift, J.H., 2012. Massive phytoplankton blooms under Arctic sea ice. *Science* 336 (1), 408.
- Arrigo, K.R., 2013. The changing Arctic Ocean. *Elem. Sci. Anth.* 1, <http://dx.doi.org/10.12952/journal.elementa.000010>.
- Bates, N.R., Mathis, T., Cooper, L.W., 2009. Ocean acidification and biologically induced seasonality of carbonate mineral saturation states in the western Arctic Ocean. *J. Geophys. Res.* 114, C11007, <http://dx.doi.org/10.1029/2008JC004862>.
- Bauerfeind, E., Nöthig, E.-M., Beszczynska, A., Fahl, K., Kaleschke, L., Kreker, K., Klages, M., Soltwedel, T., Lorenzen, C., Wegner, J., 2009. Variations in vertical particle flux in the Eastern Fram Strait (79° N/ 4° E) during 2000–2005. Results from the Deep-Sea Long-Term Observatory HAUSGARTEN. *Deep-Sea Res. I* 56, 1471–1487.
- Bauerfeind, E., Nöthig, E.-M., Pauls, B., Kraft, A., Beszczynska-Möller, A., 2014. Variability in pteropod sedimentation and corresponding aragonite flux at the Arctic deep-sea long-term observatory HAUSGARTEN in the eastern Fram Strait from 2000 to 2009. *J. Mar. Syst.* 132, L18501, 95–105, <http://dx.doi.org/10.1016/j.jmarsys.2013.12.006>.
- Belt, S.T., Massé, G., Rowland, S.J., Poulin, M., Michel, C., LeBlanc, B., 2007. A novel chemical fossil of Palaeo sea ice: IP25. *Org. Geochem.* 38, 16–27.
- Bergmann, M., Soltwedel, T., Klages, M., 2011. The interannual variability of megafaunal assemblages in the Arctic deep sea: preliminary results from the HAUSGARTEN observatory (79° N). *Deep-Sea Res.* 158, 711–723.
- Bergmann, M., Klages, M., 2012. Increase of litter at the Arctic deep-sea observatory HAUSGARTEN. *Mar. Pollut. Bull.* 64 (12), 2734–2741.
- Beszczynska-Möller, A., Fahrbach, E., Schauer, U., Hansen, E., 2012. Variability in Atlantic water temperature and transport at the entrance to the Arctic Ocean 1997–2010. *ICES J. Mar. Sci.* 69, 852–863.
- Björklund, K.R., Kruglikova, S.B., Anderson, O.R., 2012. Modern incursion of tropical Radiolaria into the Arctic Ocean. *J. Micropalaeontol.* 31, 139–158.
- Budaeva, N.E., Mokrevsky, V.O., Soltwedel, T., Gebruk, A.V., 2008. Horizontal distribution patterns in Arctic macrobenthic deep-sea communities. *Deep-Sea Res. I* 55, 1167–1178, <http://dx.doi.org/10.1016/j.dsr.2008.05.002>.
- Burrows, M.T., Schoeman, D.S., Buckley, L.B., Moore, P., Poloczanska, E.S., Brander, K.M., Brown, C., Bruno, J.F., Duarte, C.M., Halpern, B.S., Holding, J., Kappel, C.V., Kiessling, W., O'Connor, M.I., Pandolfi, J.M., Parmesan, C., Schwings, F.B., Sydeman, W.J., Richardson, A.J., 2011. The pace of shifting climate in marine and terrestrial ecosystems. *Science* 334 (6056), 652–655, <http://dx.doi.org/10.1126/science.1210288>.
- Burrows, M.T., Schoeman, D.S., Richardson, A.J., Molinos, J.C., Hoffmann, A., Buckley, L.B., Moore, P.J., Brown, C.J., Bruno, J.F., Duarte, C.M., Halpern, B.S., Hoegh-Guldberg, O., Kappel, C.V., Kiessling, W., O'Connor, M.I., Pandolfi, J.M., Parmesan, C., Sydeman, W.J., Ferrier, S., Williams, K.J., Poloczanska, E.S., 2014. Geographical limits to species-range shifts are suggested by climate velocity. *Nature* 507, 492–495, <http://dx.doi.org/10.1038/nature12976>.
- Busch, K., Bauerfeind, E., Nöthig, E.-M., 2015. Pteropod sedimentation patterns in different water depths observed with moored sediment traps over a 4-year period at the LTER station HAUSGARTEN in eastern Fram Strait. *Polar Biol.*, <http://dx.doi.org/10.1007/s00300-015-1644-9>.
- Caron, D.A., Hutchins, D.A., 2013. The effects of changing climate on microzooplankton grazing and community structure: drivers, predictions and knowledge gaps. *J. Plankton Res.* 35 (2), 235–252, <http://dx.doi.org/10.1093/plankt/fbs091>.
- Cherkasheva, A., Bracher, A., Melsheimer, C., Köberle, C., Gerdes, R., Nöthig, E.-M., Bauerfeind, E., Boetius, A., 2014. Influence of the physical environment on polar phytoplankton blooms: a case study in the Fram Strait. *J. Mar. Syst.* 132, 196–207.
- Comiso, J.C., Parkinson, C.L., Gersten, R., Stock, L., 2008. Accelerated decline in the Arctic sea ice cover. *Geophys. Res. Lett.* 35, L01703, <http://dx.doi.org/10.1029/2007GL031972>.
- de Steur, L., Hansen, E., Gerdes, R., Karcher, M., Fahrbach, E., Holfort, J., 2009. Freshwater fluxes in the East Greenland Current: a decade of observations. *Geophys. Res. Lett.* 36, L23611, <http://dx.doi.org/10.1029/2009GL041278>.
- Drinkwater, K.F., Miles, M., Medhaug, I., Ottera, O.H., Kristiansen, T., Sundby, S., Gao, Y., 2014. The Atlantic multidecadal oscillation: its manifestations and impacts with special emphasis on the Atlantic region north of 60° N. *J. Mar. Syst.* 133, 117–130.
- Duarte, C.M., Agustí, S., Wassmann, P., Arrieta, J.M., Alcaraz, M., Coello, A., Marbà, N., Hendriks, I.E., Holding, J., Garcíá-Zarandona, I., Kritzberg, E., Vaqué, D., 2012. Tipping elements in the arctic marine ecosystem. *Ambio* 41, 44–55, <http://dx.doi.org/10.1007/s13280-011-0224-7>.
- Engelsen, O., Hegseth, E.N., Hop, H., Hansen, E., Falk-Petersen, S., 2002. Spatial variability of chlorophyll-a in the Marginal Ice Zone of the Barents Sea, with relations to sea ice and oceanographic conditions. *J. Mar. Syst.* 35, 79–97.
- Fahl, K., Nöthig, E.-M., 2007. Lithogenic and biogenic particle fluxes on the Lomonosov Ridge (central Arctic Ocean) and their relevance for sediment accumulation: vertical vs. lateral transport. *Deep-Sea Res. I* 54 (8), 1256–1272.
- Fahl, K., Stein, R., 2012. Modern seasonal variability and deglacial/Holocene change of central Arctic Ocean sea-ice cover: new insights from biomarker proxy records. *Earth Planet. Sci. Lett.* 351–352, 123–133.
- Flach, E., Vanaverbeke, J., Heip, C., 1999. The meiofauna:macrofauna ratio across the continental slope of the Goban Spur (north-east Atlantic). *J. Mar. Biol. Assoc. U.K.* 79, 233–241.
- Findlay, R.H., King, G.M., Watling, L., 1989. Efficiency of phospholipid analysis in determining microbial biomass in sediments. *Appl. Environ. Microbiol.* 55, 2888–2893.
- Forest, A., Wassmann, P., Slagstad, D., Bauerfeind, E., Nöthig, E.-M., Klages, M., 2010. Relationships between primary production and vertical particle export at the Atlantic-Arctic boundary (Fram Strait HAUSGARTEN). *Polar Biol.* 33 (12), 1733–1746.
- Galeron, J., Sibuet, M., Mahaut, M.-L., Dinet, A., 2000. Variation in structure and biomass of the benthic communities at three contrasting sites in the tropical Northeast Atlantic. *Mar. Ecol. Prog. Ser.* 197, 121–137.
- Gallucci, F., Fonseca, G., Soltwedel, T., 2008. Effects of megafauna exclusion on nematode assemblages at a deep-sea site. *Deep-Sea Res. I* 55 (3), 332–349.
- Glover, A.G., Gooday, A.J., Baily, D.M., Billett, D.S.M., Chevaldonné, P., Colaco, A., Copley, J., Cuvelier, D., Desbruyères, D., Kalogeropoulou, V., Klages, M., Lampadariou, N., Lejeusne, C., Mestre, N.C., Paterson, G.L.J., Perez, T., Ruhl, H., Sarrazin, J., Soltwedel, T., Soto, E.H., Thatje, S., Tslelipidis, A., Van Gaever, S., Vanreusel, A., 2010. Temporal change in deep sea benthic ecosystems: a review of the evidence from recent time series studies. *Adv. Mar. Biol.* 58, 1–95.
- Greiser, N., Faubel, A., 1988. Biotic factors. In: Higgins, R.P., Thiel, H. (Eds.), *Introduction to the Study of Meiofauna*. Smithsonian Institution Press, Washington D.C./London, pp. 79–114.
- Grzelak, K., (PhD thesis) 2015. Structural and functional diversity of Nematoda at the Arctic deep-sea long-term observatory HAUSGARTEN (Fram Strait). Institute of Oceanology Polish Academy of Sciences, Sopot, 206 pp.
- Guilini, K., Soltwedel, T., van Oevelen, D., Vanreusel, A., 2011. Deep-sea nematodes actively colonise sediments, irrespective of the presence of a pulse of organic matter: results from an in situ experiment. *PLoS ONE* 6 (4), e18912.
- Harbison, G.R., Gilmer, R., 1986. Effects of animal behavior on sediment trap collections: implications for the calculation of aragonite fluxes. *Deep-Sea Res. Part A* 33 (8), 1017–1024.
- Hasemann, C., Bergmann, M., Kanzog, C., Loehnhofer, N., Sauter, E., Schewe, I., Soltwedel, T., 2013. Effects of dropstone induced habitat heterogeneity on Arctic deep-sea benthos with special reference to nematode communities. *Mar. Biol. Res.* 9 (3), 276–292, <http://dx.doi.org/10.1080/17451000.2012.739694>.
- Hassol, S.J., 2004. *ACIA Impacts of a Warming Arctic: Arctic Climate Impact Assessment*. Cambridge University Press.
- Hirche, H.J., Kosobokova, K., 2007. Distribution of *Calanus finmarchicus* in the northern North Atlantic and Arctic Ocean – expatriation and potential colonization. *Deep-Sea Res. II*, 2729–2747.
- Holm-Hansen, O., Lorenzen, C.J., Holmes, R.W., Strickland, J.D.H., 1965. Fluorometric determination of chlorophyll. *J. Cons. Perm. Int. Pour l'Explor. Mer* 30, 3–15.
- Hop, H., Falk-Petersen, S., Svendsen, H., Kwasniewski, S., Pavlov, V., Pavlova, O., Sørensen, J.E., 2006. Physical and biological characteristics of the pelagic system across Fram Strait to Kongsfjorden. *Prog. Oceanogr.* 71, 182–231.
- Hoste, E., (PhD thesis) 2007. Temporal and spatial variability in deep-sea meiobenthic communities from the Arctic Marginal Ice Zone. University of Gent, 201 pp.
- Hoste, E., Vanhove, S., Schewe, I., Soltwedel, T., Vanreusel, A., 2007. Spatial and temporal variations in deep-sea meiobenthic assemblages in the Marginal Ice Zone of the Arctic Ocean. *Deep-Sea Res. I* 54 (1), 109–129, <http://dx.doi.org/10.1016/j.dsr.2006.09.007>.
- Huntington, H., Boyle, M., Flowers, G., Weatherly, J., Hinzman, L., Zulueta, R., Nicolson, C., Overpeck, J., 2007. The influence of human activity in the Arctic on climate and climate impacts. *Clim. Change* 82, 77–92.
- IPCC, 2013. Climate change 2013: the physical science basis. In: Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M. (Eds.), *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom/New York, NY, USA, 1535 pp.
- Jacob, M., Soltwedel, T., Boetius, A., Ramette, A., 2013. Biogeography of deep-sea benthic bacteria at regional scale (LTER HAUSGARTEN, Fram Strait, Arctic). *PLoS ONE* 8 (9), e72779.
- Jacob, M., (PhD thesis) 2014. Influence of Global Change on microbial communities in Arctic sediments. University of Bremen, 178 pp.
- Jones, D.O.B., Yool, A., Wei, C.-L., Henson, S.A., Ruhl, H.A., Watson, R.A., Gehlen, M., 2013. Global reductions in seafloor biomass in response to climate change. *Glob. Change Biol.* 20 (6), 1861–1872, <http://dx.doi.org/10.1111/gcb.12480>.
- Kahru, M., Brota, V., Manzano-Sarabia, M., Mitchell, B.G., 2010. Are phytoplankton blooms occurring earlier in the Arctic? *Glob. Change Biol.* 17 (4), 1733–1739.
- Kanzog, C., Ramette, A., Quéréc, N.V., Klages, M., 2009. Response of benthic microbial communities to chitin enrichment: an in situ study in the deep Arctic Ocean. *Polar Biol.* 32, 105–112.
- Kortsch, S., Primicerio, R., Beuchel, F., Renaud, P.E., Rodriguez, J., Lønne, O.J., Gulliksen, B., 2012. Climate-driven regime shifts in Arctic marine benthos. *Proc. Natl. Acad. Sci. U. S. A.* 109 (35), 14052–14057.
- Kraft, A., Bauerfeind, E., Nöthig, E.-M., 2011. Amphipod abundance in sediment trap samples at the long-term observatory HAUSGARTEN (Fram Strait ~79° N/4° E). Variability in species community patterns. *Mar. Biodiv.* 41, 353–364, <http://dx.doi.org/10.1007/s12526-010-0052-1>.

- Kraft, A., Bauerfeind, E., Nöthig, E.-M., Bathmann, U., 2012. Size structure and life cycle patterns of dominant pelagic amphipods collected as swimmers in sediment traps in the eastern Fram Strait. *J. Mar. Syst.* 95, 1–15, <http://dx.doi.org/10.1016/j.jmarsys.2011.12.006>.
- Kraft, A., Nöthig, E.-M., Bauerfeind, E., Wildish, D.J., Pohle, G.W., Bathmann, U., Beszczynska-Möller, A., Klages, M., 2013. First evidence of reproductive success in a southern invader indicates possible community shifts among Arctic zooplankton. *Mar. Ecol. Prog. Ser.* 493, 291–296, <http://dx.doi.org/10.3354/meps10507>.
- Kremling, K., Lentz, U., Zeitzschel, B., Schultz-Bull, D.E., Duinker, J.C., 1996. New type of time-series sediment trap for the reliable collection of inorganic and organic trace chemical substances. *Rev. Sci. Instrum.* 67, 4360–4363.
- Kröncke, I., 1994. Macrobenthos composition, abundance and biomass in the Arctic Ocean along a transect between Svalbard and the Makarov Basin. *Polar Biol.* 14, 14519–14529.
- Kröncke, I., 1998. Macrofauna communities in the Amundsen Basin, at the Morris Jesup Rise and at the Yermak Plateau (Eurasian Arctic Ocean). *Polar Biol.* 19, 383–392.
- Kwok, R., Rothrock, D.A., 2009. Decline in Arctic sea ice thickness from submarine and ICESat records: 1958–2008. *Geophys. Res. Lett.* 36, L15501.
- Lalande, C., Bauerfeind, E., Nöthig, E.-M., Beszczynska-Möller, A., 2013. Impact of a warm anomaly on export fluxes of biogenic matter in the eastern Fram Strait. *Prog. Oceanogr.* 109, 70–77.
- Laxon, S.W., Giles, K.A., Ridout, A.L., Wingham, D.J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., Hendricks, S., Krishfield, R., Kurtz, N., Farrell, S., Davidson, M., 2013. CryoSat-2 estimates of Arctic sea ice thickness and volume. *Geophys. Res. Lett.* 40 (4), 732–737.
- Lee, C., Wakeham, S., Hedges, J.S., 1989. Sediment trap measurements of particle fluxes: effects of bacteria and zooplankton. In: GOFS (Ed.), *Sediment trap technology. Report of the U.S. GOFS Working Group on Sediment Trap Technology and Sampling*. US GOFS Planning and Coordination Office, Woods Hole, pp. 67–68.
- Lenton, T.M., Held, H., Kriegler, E., Hall, J.W., Lucht, W., Rahmsdorf, S., Schellnhuber, H.J., 2008. Tipping elements in Earth's climate system. *Proc. Natl. Acad. Sci. U. S. A.* 105, 1786–1793.
- Li, W.K.W., McLaughlin, F.A., Lovejoy, C., Carmack, E.C., 2009. Smallest algae thrive as the Arctic Ocean freshens. *Science* 326, 539.
- Lischka, S., Riebesell, U., 2012. Synergistic effects of ocean acidification and warming on overwintering pteropods in the Arctic. *Glob. Change Biol.* 18 (12), 3517–3528, <http://dx.doi.org/10.1111/gcb.12020>.
- Marsh, T.L., 1999. Terminal restriction fragment length polymorphism (T-RFLP): an emerging method for characterizing diversity among homologous populations of amplification products. *Curr. Opin. Microbiol.* 2 (3), 323–327.
- Mebrahtom Kidane, Y., (Master Thesis) 2011. Distribution of unicellular plankton organisms at the AWI HAUSGARTEN (79°N/4°E) during summer in relation to a changing Arctic environment. University Bremen, ERASMUS MUNDUS Master PRG EMBC, 58 pp.
- Meyer, K., Bergmann, M., Soltwedel, T., 2013. Interannual variation in the epibenthic megafauna at the shallowest station of the HAUSGARTEN observatory (79°N, 6°E). *Biogeosciences* 10, 3479–3492.
- Morán, X.A.G.F., López-Urrutia, A., Calvo-Díaz, Li, W.K.W., 2009. Increasing importance of small phytoplankton in a warmer ocean. *Glob. Change Biol.* 16, 1137–1144.
- Müller, J., Massé, G., Stein, R., Belt, S., 2009. Extreme variations in sea ice cover for Fram Strait during the past 30 ka. *Nat. Geosci.*, <http://dx.doi.org/10.1038/NGEO665>.
- Nicolais, M., Kattlein, C., Maslanik, J., Hendricks, S., 2012. Changes in Arctic sea ice result in increasing light transmittance and absorption. *Geophys. Res. Lett.* 39, L24501, <http://dx.doi.org/10.1029/2012GL053738>.
- Obbard, R.W., Sadri, S., Wong, Y.Q., Khitun, A.A., Baker, I., Thompson, R.C., 2014. Global warming releases microplastic legacy frozen in Arctic Sea ice. *Earth's Future* 2, 2014EF000240.
- Paquette, R.G., Bourke, R.H., Newton, J.F., Perdue, W.F., 2012. The East Greenland Polar Front in autumn. *J. Geophys. Res.* 90 (C3), 4866–4882, <http://dx.doi.org/10.1029/JC090iC03p04866>.
- Parkinson, C.L., Cavalieri, D.J., Gloersen, P., Zwally, H.J., Comiso, J.C., 1999. Arctic sea ice extents, areas, and trends, 1978–1996. *J. Geophys. Res.* 104 (C9), 20837–20856.
- Peterson, B.J., Holmes, R.M., McClelland, J.W., Vorosmarty, C.J., Lammers, R.B., Shiklomanov, A.I., Shiklomanov, I.A., Rahmsdorf, S., 2002. Increasing river discharge to the Arctic Ocean. *Science* 298, 2171–2173.
- Poloczanska, E.S., Brown, C.J., Sydeman, W.J., Kiessling, W., Schoeman, D.S., Moore, P.J., Brander, K., Bruno, J.F., Buckley, L.B., Burrows, M.T., Duarte, C.M., Halpern, B.S., Holding, J., Kappel, C.V., O'Connor, M.I., Pandolfi, J.M., Parmesan, C., Schwing, F., Thompson, S.A., Richardson, A.J., 2013. Global imprint of climate change on marine life. *Nat. Clim. Change* 3, 919–925, <http://dx.doi.org/10.1038/nclimate1958>.
- Premke, K., Klages, M., Arntz, W.E., 2006. Aggregations of Arctic deep-sea scavengers at large food falls: temporal distribution, consumption rates and population structure. *Mar. Ecol. Prog. Ser.* 325, 121–135.
- Quéric, N.-V., Soltwedel, T., Arntz, W.E., 2004. Application of a rapid direct viable count method to deep-sea sediment bacteria. *J. Microbiol. Methods* 57, 351–367.
- Ramseier, R.O., Garrity, C., Bauerfeind, E., Peinert, R., 1999. Sea-ice impact on long-term particle flux in the Greenland Sea's Is Odden-Nordbukta region 1985–1996. *J. Geophys. Res.* 104 (C3), 5329–5343.
- Renner, A.H.H., Gerland, S., Haas, C., Sprreen, G., Beckers, J.F., Hansen, E., Nicolais, M., Goodwin, H., 2014. Evidence of Arctic sea ice thinning from direct observations. *Geophys. Res. Lett.* 41, 5029–5036, <http://dx.doi.org/10.1002/2014GL060369>.
- Ruhl, H., 2007. Abundance and size distribution dynamics of abyssal epibenthic megafauna in the Northeast Pacific. *Ecology* 88, 1250–1262.
- Sakshaug, E., Skjoldal, H.R., 1989. Life at the ice edge. *Ambio* 18, 60–67.
- Serreze, M., Holland, M.M., Stroeve, J., 2007. Perspectives on the Arctic's shrinking sea-ice cover. *Science* 315, 1533–1536.
- Seiler, D., Brandt, A., 1997. Seasonal occurrence of plankton Crustacea in sediment trap samples at three depth horizons in the Greenland Sea. *Polar Biol.* 17, 337–349.
- Slagstad, D., Ellingsen, I.H., Wassmann, P., 2011. Evaluating primary and secondary production in an Arctic Ocean void of summer sea ice: an experimental simulation approach. *Prog. Oceanogr.* 90 (1–4), 117–131.
- Soetaert, K., Heip, C., 1995. Nematode assemblages of deep-sea and shelf break sites in the North Atlantic and Mediterranean Sea. *Mar. Ecol. Prog. Ser.* 125 (1–3), 171–183.
- Soltwedel, T., Bauerfeind, E., Bergmann, M., Budaeva, N., Hoste, E., Jaeckisch, N., Juterzenka, K.v., Matthiesen, J., Mokievsky, V., Nöthig, E.-M., Quéric, N., Sablotny, B., Sauter, E., Schewe, I., Urban-Malinga, B., Wegner, J., Włodarska-Kowalcuk, M., Klages, M., 2005. HAUSGARTEN: multidisciplinary investigations at a deep-sea, long-term observatory in the Arctic Ocean. *Oceanography* 18 (3), 46–61.
- Soltwedel, T., Jaeckisch, N., Ritter, N., Hasemann, C., Bergmann, M., Klages, M., 2009. Bathymetric patterns of megafaunal assemblages from the arctic deep sea observatory HAUSGARTEN. *Deep-Sea Res. I* 56 (10), 1856–1872, <http://dx.doi.org/10.1016/j.dsrs.2009.05.012>.
- Soltwedel, T., Volkenandt, M., Mokievsky, V., Hasemann, C., Sauter, E., Rabouille, C., 2013. Effects of experimentally increased near-bottom flow on meiofauna diversity in the deep Arctic Ocean. *Deep-Sea Res.* 173, 31–45.
- Steinacher, M., Joos, F., Frölicher, T.L., Plattner, G.-K., Doney, S.C., 2009. Imminent ocean acidification in the Arctic projected with the NCAR global coupled carbon cycle-climate model. *Biogeosciences* 6, 515–533.
- Thiel, H., 1978. Benthos in upwelling regions. In: Boje, R., Tomczak, M. (Eds.), *Upwelling Ecosystems*. Springer Verlag, Berlin, pp. 124–138.
- Tremblay, J.-É., Gagnon, J., 2009. The effects of irradiance and nutrient supply on the productivity of Arctic waters: a perspective on climate change. In: Nihoul, J.C.J., Kostianoy, A.G. (Eds.), *Influence of Climate Change on the Changing Arctic and Sub-Arctic Conditions*. Springer, The Netherlands, pp. 73–93.
- v. Bodungen, B., Wunsch, M., Fürderer, H., 1991. Sampling and analysis of suspended particles in the northern north Atlantic. In: Hurd, D.C., Spencer, D.W. (Eds.), *Marine Particles and Characterization*, vol. 63. *Geophys. Monogr.*, pp. 47–56.
- Wang, M., Overland, J.E., 2012. A sea ice free summer Arctic within 30 years: an update from CMIP5 models. *Geophys. Res. Lett.* 39, <http://dx.doi.org/10.1029/2012GL052868>.
- Wassmann, P., Slagstad, D., Ellingsen, I., 2010. Primary production and climatic variability in the European sector of the Arctic Ocean prior to 2007: preliminary results. *Polar Biol.* 33 (12), 1641–1650.
- Wassmann, P., Duarte, C.M., Agusti, S., Sejr, M.K., 2011. Footprints of climate change in the Arctic marine ecosystem. *Glob. Change Biol.* 17, 1235–1249.
- Weydmann, A., Carstensen, J., Gosczko, I., Dmoch, K., Olszewska, A., Kwasniewski, S., 2014. Shift towards the dominance of boreal species in the Arctic: inter-annual and spatial zooplankton variability in the West Spitsbergen Current. *Mar. Ecol. Prog. Ser.* 501, 41–52, <http://dx.doi.org/10.3354/meps10694>.
- Willis, K.J., Cottier, F.R., Kwaśniewski, S., 2008. Impact of warm advection on the winter zooplankton community in an Arctic fjord. *Polar Biol.* 31, 475–481.
- Włodarska-Kowalcuk, M., Kendall, M.A., Węsławski, J.-M., Klages, M., Soltwedel, T., 2004. Depth gradients of benthic standing stock and on the continental margin at a high latitude ice-free site (off West Spitsbergen 79 N). *Deep-Sea Res. I* 51, 1903–1914.
- Wohlers, J., Engel, A., Zöllner, E., Breithaupt, P., Jürgens, K., Hoppe, H.-G., Sommer, U., Riebesell, U., 2009. Changes in biogenic carbon flow in response to sea surface warming. *Proc. Natl. Acad. Sci. U. S. A.* 106, 7067–7072.
- Wulff, T., Lehmenhecker, S., Bauerfeind, E., Hoge, U., Shurn, K., Klages, M., 2013. Biogeochemical research with an autonomous underwater vehicle: payload structure and arctic operations. MTS/IEEE OCEANS Bergen, <http://dx.doi.org/10.1109/OCEANS-Bergen.2013.6608043>.
- Yentsch, C.S., Menzel, D.W., 1963. A method for the determination of phytoplankton chlorophyll and phaeophytin by fluorescence. *Deep-Sea Res.* 10, 221–231.
- Zhang, F., He, J., Lin, L., Jin, H., 2015. Dominance of picophytoplankton in the newly open surface water of the central Arctic Ocean. *Polar Biol.*, <http://dx.doi.org/10.1007/s00300-015-1662-7>.