Daily up and down of the plankton animals in the sea

New study on the migration behaviour of zooplankton, its diurnal and annual cycles

A unique series of measurements taken over several years in the Antarctic Ocean provide new findings about the daily vertical migration of zooplankton communities. Scientists of the Thünen Institute of Sea Fisheries in Hamburg and the Alfred Wegener Institute observed changes during the year and between years. The Antarctic zooplankton is the main source of food for many fish and whale species, including the largest mammal in the world, the blue whale.

For the first time, researchers were able to document the annual cycle of zooplankton communities in the Southern Ocean. Using data from anchored acoustic Doppler current profilers (ADCP), Dr Boris Cisewski of the Thünen Institute of Sea Fisheries and Dr Volker Strass of the AWI continuously recorded the daily vertical migration of the zooplankton for three years. They also investigated how individual environmental parameters such as light, sea ice cover, food availability, circulation and temperature as well as salinity influence their migration behaviour.

The daily vertical migration of zooplankton -- often crustaceans with body lengths ranging from millimetres to centimetres -- is mainly triggered by the day-night cycle. In order to escape potential predators, they dive into the dark depths at sunrise and stay there during the day. After sunset, they once again rise to the upper layers to feed where the sunlight has allowed planktonic algae to grow. Until now, only short time snippets of the migration pattern of the zooplankton in the Southern Ocean existed. Because of its seasonal sea ice cover, many areas are not accessible by ship during the southern winter. At this time of year, biological network samples can only be taken intermittently.

The current study is based on data that was collected during three Polarstern expeditions and with deep-sea moorings deployed in the Southern Ocean between 2005 and 2008 within the framework of the LAKRIS project (Lazarev Sea Krill Study). As part of this study ADCPs were moored at three different geographical locations along the Greenwich meridian; the ADCPs send out sound waves at fixed intervals and cover an up to 500 m deep water layer under the surface. While the strength of the echo provides information about the concentration of the zooplankton, the migration velocity can be calculated based on the Doppler shift of the sound frequency.

On the basis of these long-term measurements, the researchers examined not only daily, but seasonal vertical migration patterns of individual zooplankton communities. One of the things they found was that zooplankton discontinue their migration behaviour for a few weeks at the beginning of the southern summer from end of November up until January. It appears that the abundance of food in the surface water is so attractive for the animals that they do not recede to greater depths during the day to hide from predators. Further evidence for this hypothesis is provided by measurements of the sea ice and chlorophyll distribution. When the ice begins to melt, so-called ice-edge algal blooms are formed. They are the main source of food for Antarctic zooplankton at this time of year. Surprising were the large differences of the zooplankton abundances between the three years under study. The causes of these interannual differences have not yet been fully established. It needs more physical and biological measurements in order to estimate future changes to the ecosystem of the Southern Ocean as a result of climate change.

Journal Reference: Boris Cisewski, Volker H. Strass. Acoustic insights into the zooplankton dynamics of the eastern Weddell Sea. *Progress in Oceanography*, 2016; 144: 62 DOI: 10.1016/j.pocean.2016.03.005 pdf reprint follows......

Progress in Oceanography 144 (2016) 62-92

Contents lists available at ScienceDirect

Progress in Oceanography

journal homepage: www.elsevier.com/locate/pocean

Acoustic insights into the zooplankton dynamics of the eastern Weddell Sea



^a Thünen-Institut für Seefischerei, Palmaille 9, 22767 Hamburg, Germany ^b Alfred-Wegener-Institut, Helmholtz-Zentrum für Polar- und Meeresforschung, Am Handelshafen 12, 27570 Bremerhaven, Germany

ARTICLE INFO

Article history: Received 17 March 2015 Received in revised form 22 March 2016 Accepted 23 March 2016 Available online 1 April 2016

ABSTRACT

The success of any efforts to determine the effects of climate change on marine ecosystems depends on understanding in the first instance the natural variations, which contemporarily occur on the interannual and shorter time scales. Here we present results on the environmental controls of zooplankton distribution patterns and behaviour in the eastern Weddell Sea, Southern Ocean. Zooplankton abundance and vertical migration are derived from the mean volume backscattering strength (MVBS) and the vertical velocity measured by moored acoustic Doppler current profilers (ADCPs), which were deployed simultaneously at 64°S, 66.5°S and 69°S along the Greenwich Meridian from February, 2005, until March, 2008. While these time series span a period of full three years they resolve hourly changes.

A highly persistent behavioural pattern found at all three mooring locations is the synchronous diel vertical migration (DVM) of two distinct groups of zooplankton that migrate between a deep residence depth during daytime and a shallow depth during nighttime. The DVM was closely coupled to the astronomical daylight cycles. However, while the DVM was symmetric around local noon, the annual modulation of the DVM was clearly asymmetric around winter solstice or summer solstice, respectively, at all three mooring sites. DVM at our observation sites persisted throughout winter, even at the highest latitude exposed to the polar night. Since the magnitude as well as the relative rate of change of illumination is minimal at this time, we propose that the ultimate causes of DVM separated from the light-mediated proximal cue that coordinates it. In all three years, a marked change in the migration behaviour occurred in late spring (late October/early November), when DVM ceased. The complete suspension of DVM after early November is possibly caused by the combination of two factors: (1) increased availability of food in the surface mixed layer provided by the phytoplankton spring bloom, and (2) vanishing diurnal enhancement of the threat from visually oriented predators when the illumination is quasi-continuous during the polar and subpolar summer.

Zooplankton abundance in the water column, estimated as the mean MVBS in the depth range 50–300 m, was highest end of summer and lowest mid to end winter on the average annual cycle. However, zooplankton abundance varied several-fold between years and between locations. Based on satellite and in situ data of chlorophyll and sea ice as well as on hydrographic measurements, the interannual and spatial variations of zooplankton mean abundance can be explained by differences in the magnitude of the phytoplankton spring bloom, which develops during the seasonal sea ice retreat. Whereas the vernal ice melt appears necessary to stimulate the blooming of phytoplankton, it is not the determinator of the blooms magnitude, its areal extent and duration. A possible explanation for the limitation of the phytoplankton bloom in some years is top-down control. We hypothesise that the phytoplankton spring development can be curbed by grazing when the zooplankton had attained high abundance by growth during the preceding summer.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

E-mail address: boris.cisewski@thuenen.de (B. Cisewski).

Zooplankton constitutes the central hub of pelagic ecosystems. It ensures the flow of energy, harvested from the sun light and stored in organic substances by phytoplankton during photosynthesis, from the primary producers to the higher trophic levels







^{*} Corresponding author at: Thünen-Institut für Seefischerei, Palmaille 9, 22767 Hamburg, Germany. Tel.: +49 40 38905 224; fax: +49 40 38905 263.

and thus to the marine living resources exploitable by men. Zooplankton also plays a role in the biological pump, hence in the uptake of CO₂ from the atmosphere and the subsequent transfer of biogenic carbon to the deep ocean (Longhurst and Harrison, 1988). To the biological pump it contributes by two principal ways: first by release of sinking fecal pellets formed from the ingested organic-carbon-rich food; second by migrating vertically between feeding strata near the sun-lit surface where the phytoplankton grows and deeper resting layers, where not only defecation but also excretion of dissolved compounds of carbon and nutrient elements occurs, and in consequence accounts for a transport of matter from surface to depth (Steinberg et al., 2000; Turner, 2015). However, zooplankton may also intercept the vertical flux of organic carbon-rich particles by grazing on sinking phytoplankton aggregates and on fecal pellets (Wallace et al., 2013). If the subsequent respiration occurs relocated towards the surface, an upward flux component is created which reduces the overall downward carbon flux.

A widespread characteristic behavioural pattern of marine and fresh water zooplankton is a diel vertical migration (DVM), in which the vertical distribution of the migrating zooplankters changes over the 24-h day. Despite many decades of study, both the evolutionary mechanisms behind DVM and the cues influencing it (Lampert, 1989; Cohen and Forward, 2009) are still under debate. However, predator evasion is currently the most favoured hypothesis. The most-common, so-called nocturnal vertical migration brings migrants from typically the bathypelagic or mesopelagic zones into the epipelagic zone at night to feed, and back to depth in daytime to decrease mortality by reducing the risk of detection by visually hunting predators (Zaret and Suffern, 1976; Lampert, 1989).

DVM is considered a behavioural response to a combination of exogenous environmental factors (e.g. light, gravity, temperature, salinity, oxygen, and hydrostatic pressure) and endogenous factors (e.g. sex, age, changes in behaviour and physiology) originating from the organism itself (Forward, 1988). Daylight is generally assumed the dominant exogenous cue controlling DVM, because the times of migration usually correspond to the times of the rapid light intensity changes around sunrise and sunset (see Roe, 1974; Forward, 1988; Haney, 1988; Ringelberg, 1995). The three main hypotheses for the use of light as an exogenous stimulus for DVM are (1) preferendum or isolume hypothesis, (2) absolute intensity threshold hypothesis and (3) the rate-of-change hypothesis (see review of Cohen and Forward, 2009). The preferendum or isolume hypothesis postulates that zooplankton follow a preferred or optimal light level during vertical migration. The absolute intensity threshold hypothesis states that ascent and descent by an organism are initiated when the ambient light intensity changes below or above a specific level. Finally, the rate of change hypothesis suggests that vertical movements in DVM are caused by behavioural responses to the relative rate and direction of irradiance change from some non-fixed ambient adaption intensity.

Some authors have also addressed the effect of advection combined with DVM on the distribution (dispersal or retention) of zooplankton whereby the organisms take advantage of horizontal currents with vertically varying speeds and directions during their upward and downward movements (Hardy and Gunther, 1935; Wroblewski, 1982; Manuel and O'Dor, 1997; Eiane et al., 1998; Krägefsky, 2008; Krägefsky et al., 2009).

Driven by the annual light cycle, polar environments exhibit an extreme seasonal variability. Poleward of the latitude of the Polar Circle the sun stays below the horizon between the autumnal and vernal equinoxes and above the horizon between the spring equinox and the next autumn. For marine organisms, the change between continuous light during summer and continuous darkness during winter is additionally modified by the seasonal build-up and decay of the sea ice cover on large horizontal scale. The sea ice around Antarctica typically covers more than 20 million km² in winter and recedes to less than 4 million km² during spring and summer (Zwally et al., 1983; Gloersen and Campbell, 1991). A paradigm of marine polar biology is that the seasonal changes in ambient sun light and in the sea ice coverage have a profound influence on the dynamics of high-latitude marine ecosystem processes.

In the austral winter and spring, seasonal pack ice can contain algal concentrations that are one to three orders of magnitude higher than in the water column below and provide a crucial food source for grazers such as Antarctic krill (Euphausia superba) when the food resources in the water column are very low (Quetin and Ross, 2009). During the austral summer, when the sea ice edge recedes, low-salinity melt water usually stabilizes the water column and creates a shallow mixed-laver, which facilitates primary production in a high irradiance environment and accumulation of pelagic phytoplankton biomass, which provides the major food supply for the zooplankton during this time of the year (Smith and Nelson, 1985). The extreme variations in daylight hours, with twenty-four hours of daylight in summer, and complete darkness at mid-winter, raise the question of whether high-latitude zooplankton and nekton display DVM during the polar night and the midnight sun, when the relative change in irradiance $\Delta I/I$ is small (e.g. Buchanan and Haney, 1980; Fortier et al., 2001; Blachowiak-Samolyk et al., 2006; Cottier et al., 2006; Falk-Petersen et al., 2008; Berge et al., 2009; Benoit et al., 2010; Rabindranath et al., 2010).

Since the pioneering work of Flagg and Smith (1989), many investigators have utilised the echo intensity measured by acoustic Doppler current profilers (ADCPs) to examine the temporal and spatial variability in the distribution of zooplankton biomass and diel vertical migration for over two decades, presuming that zooplankton are the main scatterers of sound in the typically used frequency range of 10 s to 100 s kHz (e.g., Heywood et al., 1991; Fischer and Visbeck, 1993; Flagg et al., 1994; Heywood, 1996; Ashiian et al., 1998: Luo et al., 2000: Pinot and Jansá, 2001: Véle z-Belchí et al., 2002: Jiang et al., 2007: Cisewski et al., 2010: Radenac et al., 2010; Wallace et al., 2010). ADCPs can complement conventional net sampling in an excellent way because they can be deployed in extreme environments and can operate autonomously year-round, and thus provide sufficient spatial and temporal resolution to yield precise estimates of timing, velocity and extent of DVM. They also have the advantage of not being subject to avoidance problems as nets, imposed by zooplankton with sufficient swimming power to escape moving objects (Burd and Thomson, 2012). On the other hand, however, they have the drawback of reduced ability of identification of species and of measuring quantitatively numbers of individuals and biomass.

Building on time series recorded over three years by selfcontained ADCPs, moored at three different locations along the Greenwich Meridian in the eastern Weddell Sea, we here present an analysis of the meridional and interannual variability in the measured vertical velocity and in the mean volume backscattering strength (MVBS). This study thus provides an acoustic insight into the zooplankton vertical migration pattern and abundance changes in a sub-region of the barely sampled Southern Ocean. The presented time series data enable to track the changes that occur over three years at hourly resolution. The particular purposes of this study are (1) to analyse and interpret the derived DVM patterns and abundance variations for their seasonal modulation and interannual variability, (2) to scrutinise the annual asymmetry of the diel vertical migration first described by Cisewski et al. (2010) for its recurrence, and (3) to investigate how the environmental conditions control the distribution of zooplankton spatially and temporally.

Table 1Summary of cruise dates.

Cruise	Objective	Start	End	No. CTD stats.	Cruise report
ANT-XXIII/2	Summer 2005/2006 survey	11/19/2005	01/12/2006	87	Strass (2007)
ANT-XXIII/6	Winter 2006 survey	06/17/2006	08/21/2006	90	Bathmann (2008)
ANT-XXIV/2	Summer 2007/2008 survey	11/28/2007	02/03/2008	95	Bathmann (2010)

2. Materials and methods

2.1. Sampling programme and study area

Three expedition cruises were conducted with R.V. Polarstern during the years 2005-2008 in the eastern Weddell Sea, also known as the Lazarev Sea, as a contribution to the LAKRISproject (Lazarev Sea Krill Study) and to the International Polar Year. During these surveys, scheduled in order to also reveal seasonal variations, a multitude of physical and biological data were collected. The core data set analysed in this study consists of (1) measurements of the 3D velocity field and of backscatter strength recorded by self-contained acoustic Doppler current profilers (SCADCPs) moored at three different locations (64°, 66.5° and 69°S, 0°E) between February 2005 and March 2008 and (2) three interdisciplinary surveys that were conducted in early summers 2005/2006 (S05/06), 2007/2008 (S07/08) and winter 2006 (W06) (Table 1). These surveys cover the area between 6°W and 3°E and from the Antarctic continental shelf at roughly 70°S to 60°S (Fig. 1). The study area is characterised by a marked bottom topography comprising the Antarctic continental shelf, deep abyssal plains and an isolated seamount, Maud Rise. It exhibits a complex pattern of ocean circulation, which determines the inflow of Warm Deep Water (WDW) into the Weddell Sea (Bersch et al., 1992; Gordon and Huber, 1995; Muench et al., 2001; de Steur et al., 2007; Leach et al., 2011; Cisewski et al., 2011).

2.1.1. Hydrographic data

Between November 2005 and February 2008, 272 casts with a CTD type Sea-Bird Electronics SBE 911plus were made during the three LAKRIS surveys, in order to map the hydrographic fields of the study area. Of these, 106 extended to full ocean depth, while the others were limited mostly to the upper 1000 m of the water column. All CTD stations were organised on a regular grid, made up of four meridional sections running between 60°S and 70°S and along 3°E, 0°E, 3°W and 6°W during the summer cruise 2005/2006. However, the entire transect along 6°W of the planned grid had to be abandoned due to difficult ice conditions and logistical constraints during the other two surveys in winter 2006 and summer 2007/2008. The distance between stations along the meridional sections was nominally 30 nautical miles. Water samples were collected with a Sea-Bird Carousel sampler with 24 12-l bottles. For in situ calibration, temperatures were measured with a digital thermometer Sea-Bird SBE35. The conductivity readings of the CTD probe were corrected by means of the salinities of water samples, which were determined on board by a Guildline Autosal 8400A salinometer referenced to IAPSO standard seawater. The temperature sensor was calibrated by the manufacturer a few months prior to the cruise and afterwards to an accuracy better than 0.001 K for all surveys. Salinity derived from the CTD measurements was calibrated to a final accuracy of better than 0.002 for all cruises by comparison to the salinity samples.

2.1.2. Vessel-mounted acoustic Doppler current profiler

During the listed surveys (Table 1) current velocities were measured continuously along the ship's track down to 335 m depth using a vessel-mounted acoustic Doppler current profiler (VMADCP type Ocean Surveyor; Teledyne RD Instruments USA, Poway, California, 153.6 kHz nominal frequency). The transducers were located 11 m below the water line and were protected against ice floes by an acoustically transparent plastic window. The velocity components u (eastward), v (northward) and w (upward) were averaged in 2 min ensembles in 4 m thick depth bins between 19 and 335 m depth. The ADCP processing was carried out by using the CODAS3 software package developed by E. Firing and colleagues (Firing, 1991). Barotropic tidal currents were calculated using the Circum-Antarctic Tide Model CATS02.01 developed by Padman et al. (2002) and subsequently removed. For further information we refer to Cisewski et al. (2011).

2.1.3. Self-contained acoustic Doppler current profilers

The self-contained SCADCPs, deployed as components of oceanographic moorings, were of type Workhorse Longranger (Teledyne RD Instruments USA, Poway, California) using a fourbeam, convex configuration with a beam angle of 20° and a frequency of 76.8 kHz. They were moored at nominal depths between 309 and 477 m (Table 2) in upward-looking mode and measured horizontal and vertical currents and acoustic backscatter intensity from roughly 20 to 480 m. The SCADCPs were employed in three different configurations depending on the instrument's internal data storage capacity (Table 3). While the SCADCP data of the first deployment (early February to mid of December 2005) were analysed and presented by Cisewski et al. (2010), we focus here on the second deployment period continuing from December 2005 to early March 2008. For the latter, the number of depth cells was set to 80 with a bin size of 8 m. The sampling interval was set to one ping per ensemble with a ping rate of one ping every 2-10 min (Table 3). The moorings were deployed in December 2005 during R.V. Polarstern cruise ANT-XXIII/2 and recovered during R.V. Polarstern cruise ANT-XXIV/3 (Table 2).

2.1.4. Other environmental parameters

In order to examine for environmental factors that possibly exert a control on zooplankton behaviour and abundance, various variables were considered. Sunrise/sunset times and sun angles at the mooring locations were calculated with the solar position algorithm (SPA) introduced by Reda and Andreas (2004). The sea ice coverage and thickness were estimated from four upward looking sonars (ULS), which were placed at about 100-150 m depth below the sea surface at the top of the oceanographic moorings. The ULS data processing technique is described by Strass (1998) with modifications introduced by Behrendt et al. (2013). Because of data gaps occurring during the ULS measurements we also use Special Scanning Microwave Imager (SSM/I)-derived sea ice concentration data, which are provided by the National Snow and Ice Center, Boulder, Colorado. For this study we estimated the sea ice coverage at 64°S, 66.5°S and 69°S from daily means of the available ULS data and compared them to the nearest pixel of SSM/I data derived from the Bootstrap algorithm. Bootstrap algorithm data (Comiso, 1999) were retrieved from NSIDC via http://www. nsidc.org/data/nsidc-0079.html. It has to be noted that the ULS footprint is approximately 10 m, whereas the SSM/I data is provided on a 25-km grid.

The concentration of chlorophyll a was measured at every CTD station during R.V. *Polarstern* cruises ANT-XXIII/2 (Fach et al., 2007), ANT-XXIII/6 (Herrmann, 2009) and ANT-XXIV/2



Fig. 1. Bathymetric map of the Atlantic and Indian Ocean sectors of the Southern Ocean including the study area.

 Table 2

 Data coverage of the acoustic Doppler current profilers (ADCP) time series.

Mooring	Position	Water depth (m)	Mean ADCP depth	Time period	Duration (d)
AWI-229/6 ^a	63°57.16′S 00°00.37′W	5200	346.9 ± 56.7 m	Feb. 7, 2005-Dec. 16, 2005	313
AWI-229/7	63°57.17′S 00°00.17′W	5200	389.1 ± 25.7 m	Dec. 16, 2005-Feb. 28, 2008	805
AWI-231/6 ^a	66°30.66′S 00°01.91′W	4540	317.1 ± 0.8 m	Feb. 9, 2005–Dec. 18, 2005	313
AWI-231/7	66°30.67′S 00°01.90′W	4540	309.0 ± 1.6 m	Dec. 18, 2005-Mar. 6, 2008	810
AWI-232/7 ^a	68°59.75′S 00°00.11′W	3370	379.2 ± 3.0 m	Feb. 17, 2005-Dec. 19, 2005	306
AWI-232/8	68°59.75'S 00°00.16'W	3370	477.3 ± 2.5 m	Dec. 19, 2005-Mar. 10, 2008	813

^a For further details, see Cisewski et al. (2010).

Table 3

Instrument configurations	for the self-contained	acoustic Dopple	er current profilers	(ADCP)
---------------------------	------------------------	-----------------	----------------------	--------

Mooring	Frequency (kHz)	Bin length (m)	Number of bins	Sampling rate	Sampling interval (min)
AWI-229/6 ^a	76.8	16	38	1	10
AWI-229/7	76.8	8	80	1	4
AWI-231/6 ^a	76.8	8	80	1	2
AWI-231/7	76.8	8	80	1	4
AWI-232/7ª	76.8	16	38	1	10
AWI-232/8	76.8	8	80	1	4

^a For further details, see Cisewski et al. (2010).

(Bathmann, 2010). These discrete in situ chlorophyll data was supplemented by maps of the near-surface chlorophyll concentration derived from satellite remote sensing. Satellite-sensed chlorophyll distributions were acquired using products processed and distributed by ACRI-ST GlobColour service, supported by EU FP7 MyOcean and SA GlobColour Projects, using ESA ENVISAT Medium Resolution Imaging Spectrometer (MERIS) data, NASA Moderate Resolution Imaging Spectrometer (MODIS) and Sea-Viewing Wide Field of View Sensor (SeaWiFS) data. The obtained maps indicate the monthly mean surface concentrations within the study area for the months January, February, March, November and December of the consecutive years 2005–2008. Furthermore, based on eightday composites, we estimated time series of the surface chlorophyll concentration at the three mooring locations; we used a grid of 3×3 pixels around the mooring location and computed the average. The GlobColour products have been derived with the GSM model and algorithm, developed by Maritorena and Siegel (2005).

2.2. Data analysis

2.2.1. MVBS computation

The Workhorse Longranger ADCPs recorded echo intensity on a 0-255 automatic gain control (AGC) count scale. The echo intensity *E* was converted to the mean volume backscattering strength

MVBS (dB) after the version of the sonar equation presented by Deines (1999):

 $MVBS = C + 10\log_{10}((T_x + 273.16)R^2) - L_{DBM} - P_{DBW} + 2\alpha R + K_C(E - E_r))$

where C is a system constant delivered by the manufacturer (which includes transducer and system noise characteristics and is -159.1 dB for the Workhorse Longranger), L_{DBM} is the $10\log_{10}$ (transmit pulse length/meter), P_{DBM} is the $10\log_{10}$ (transmit power/Watt), T_x is the temperature of the transducer (°C), R is the range along the beam to scatterers (m), α is the sound absorption coefficient of seawater (dB/m) and K_c is a beam-specific scaling factor (dB/count). We followed Brierley et al. (2006) and used beam-averaged data because the four beams together gave a better signal-to-noise ratio than individual beams. The noise level (E_r) of all four beams was determined from the minimum values of RSSI (Received Signal Strength Indicator) counts obtained in the remotest depth cell, when the sea surface was outside the ADCP range. Sound velocity *c* and sound absorption coefficient α were considered variable with depth and time and calculated according to the UNESCO formula introduced by Fofonoff and Millard (1983) and computed after Francois and Garrison (1982). For determination of the vertical distribution of both parameters in the depth range above the ADCP, we used CTD profiles measured at the mooring positions during deployment and recovery. In order to construct a seasonal cycle of the sound velocity and absorption for the full deployment period between February 2005 and March 2008, we interpolated in time all available and occasional CTD profiles, which were conducted at the mooring sites.

2.2.2. Estimation of the vertical (migration) velocity

In order to analyse the seasonal variability of the vertical (migration) velocity, mean diel cycles of the ADCP-derived vertical velocity were calculated for successive months. Since the velocity uncertainty of single-ping ADCP data is too large, averaging is used to reduce the measurement uncertainty to acceptable levels. The manufacturer indicates a single-ping uncertainty of 146 mm s⁻¹ for their instruments configured in longrange mode with a bin length of 8 m. Because individual pings are independent, the statistical uncertainty of the measurements can be reduced according the equation:

 $\frac{\text{Statistical uncertainty for one ping}}{\sqrt{\text{Number of Pings}}}$



Fig. 2. Bathymetry, circulation and hydrography in the LAKRIS study area during R.V. *Polarstern* cruise ANT-XXIII/2 (redrawn after Cisewski et al., 2011). Tidally corrected ADCP velocity vectors are spatially averaged over 150–200 m depth and in 5 km segments along the track (a), vertical distribution of temperature (b), salinity (c), and density along the prime meridian during the period 2005/12/14 to 2005/12/20 (d). Moorings and ADCP positions are illustrated as orange squares in (a) and as white dashed lines and white squares in (b) through (d), respectively. ASF: Antarctic Slope Front; ASW: Antarctic Surface Water; WDW: Warm Deep Water.



Fig. 3. Bathymetry, circulation and hydrography in the LAKRIS study area during R.V. *Polarstern* cruise ANT-XXIII/6 (redrawn after Cisewski et al., 2011). Tidally corrected ADCP velocity vectors are spatially averaged over 150–200 m depth and in 5 km segments along the track (a), vertical distribution of temperature (b), salinity (c), and density along the prime meridian during the period 2006/06/26 to 2006/08/12 (d). Moorings and ADCP positions are illustrated as orange squares in (a) and as white dashed lines and white squares in (b) through (d), respectively. Since the CTD was exposed to deck temperatures below freezing, ice formed inside the conductivity cell at stations 500–501. Thus, measurements were low of correct until the ice layer melted and disappeared. These data were excluded from analysis. ASF: Antarctic Slope Front; ASW: Antarctic Surface Water; WDW: Warm Deep Water.

The statistical uncertainty of the velocity measurements reduces to standard errors of $(146/(\sqrt{30} \cdot \sqrt{24} \cdot \sqrt{15}) \text{ mm s}^{-1}) = 1.4 \text{ mm s}^{-1}$ if we average over 10.800 single-ping ensembles, respectively, which corresponds to the theoretical precision of *w* averaged over a month. According to the method first introduced by Plueddemann and Pinkel (1989) and modified by Luo et al. (2000) and Cisewski et al. (2010), we also infer the migration velocity for the scattering layers from the "slope" velocity $\Delta z/\Delta t$ of individual MVBS contours.

3. Results

3.1. Hydrography

-6000

-5000

-4000

-3000

Depth [m]

-2000

-1000

0

For the two summer surveys, the hydrographic conditions of the upper 500 m in the study area are illustrated by sections along the prime meridian from 60°S to 70°S (Figs. 2b–d and 4b–d). The uppermost ocean surface layer consists mainly of less saline and relatively warm Antarctic Surface Water (AASW), which caps the

subsurface temperature minimum or Winter Water (WW) layer and limits surface mixing to a mixed layer depth of 4-46 m in the latitude range north of 69°S, i.e. north of the continental shelf break. For this study, the mixed layer depth was calculated, in accordance to the definition of Cisewski et al. (2005), as the depth at which the in situ density increased by $\Delta \sigma_T = 0.02 \text{ kg m}^{-3}$ compared to the uppermost valid value of a given CTD profile. The WW is the remnant of the previous winter mixed layer stratified towards the surface by seasonal warming and freshening (Park et al., 1998). Underneath the surface layer, the Warm Deep Water (WDW; Carmack and Foster, 1975), characterised by an intermediary temperature maximum of about 1.1 °C and a salinity maximum of about 34.7. is found. In the south, the surface mixed laver of about 100-120 m depth deepens towards the shelf break to more than 600 m near the Antarctic Slope Front (Jacobs, 1991), which separates the WW and the WDW from the colder and less saline shelf waters near the Antarctic continent (Fahrbach et al., 2004). During austral winter 2006 (Fig. 3b-d), the mixed layer extended down to between 100 and 200 m, coincident with the base of the WW. The WDW layer below however reveals an almost similar

latitude [°S]



Fig. 4. Bathymetry, circulation and hydrography in the LAKRIS study area during R.V. *Polarstern* cruise ANT-XXIV/2. Tidally corrected ADCP velocity vectors are spatially averaged over 150–200 m depth and in 5 km segments along the track (a), vertical distribution of temperature (b), salinity (c), and density along the prime meridian during the period 2008/01/01 to 2008/01/06 (d). Moorings and ADCP positions are illustrated as orange squares in (a) and as white dashed lines and orange squares in (b) through (d), respectively. ASF: Antarctic Slope Front; ASW: Antarctic Surface Water; WDW: Warm Deep Water.

pattern as observed during the foregoing summer. Considered together with the temperature time series obtained from the ADCP thermistors, the upper 500 m temperature sections along 0°E (Figs. 2b, 3b and 4b) reveal that the instruments were located within the WDW inflow for the whole deployment period (February 2005–March 2008). For a more detailed description of the hydrography, we refer to Cisewski et al. (2011).

3.2. Circulation

3.2.1. Underway current observations

Figs. 2a and 4a illustrate the horizontal current vectors in the depth range 150–200 m for the summer surveys 2005/2006 (S05/06) and 2007/2008 (S07/08), measured with the VMADCP along the ship track and averaged in 5 km segments after the correction for tides. The vectors show the highest velocity within the westward flowing Antarctic Coastal Current (AntCC), which is confined to the Antarctic continental shelf slope and is associated with maximum velocities of about $21-42 \text{ cm s}^{-1}$ (S05/06) and 15–28 cm s⁻¹ (S07/08). Apart from the westward flowing AntCC, only one other persistent strong current was found in the study area, namely a westward then southwestward circulation around

the northern and northwestern flanks of Maud Rise with maximum velocities of about 20-23 cm s⁻¹ (S05/06) and 13-28 cm s⁻¹ (S07/08), respectively. In both summers, this jet is 30-60 km wide and follows the 4000 and 5000 m-isobaths, which locally correspond to the deepest part of the slope of the seamount. Above Maud Rise the water column appears rather stagnant, evidencing the Taylor column that is caused by this topographic structure and veers the flow around it (Ou, 1991; Brandt et al., 2011). Further south, between Maud Rise and the AntCC, weak recirculation occurs in narrow bands with maximum velocities of about $5-14 \text{ cm s}^{-1}$ (S05/06) and $5-19 \text{ cm s}^{-1}$ (S07/08) in varying directions. At the two westernmost transects the velocity field is dominated by a number of mesoscale features indicative of eddies. Fig. 3a reveals the horizontal currents in the depth range 150–200 for the winter survey 2006 (W06); along the 6°W transect and at the southernmost stations of the 0°E and 3°E transects no measurements were possible because of severe ice conditions. Unfortunately, there is also a gap in the VMADCP data due to a failure in the ship's navigational system along the 3°E transect between 63.5°S and 64°S, which coincides with a major part of the northern gateway for WDW. The vectors along the 3°E transect reveal the highest velocities within the AntCC of about 34 cm s⁻¹. However,



Fig. 5. Time series of daily mean horizontal current speeds measured by the moored ADCPs, spatially averaged between 100 and 200 m depth, and corresponding current frequency roses showing the distribution of current directions and speeds at 64°S (a), 66.5°S (b), and 69°S (c) over the full deployment period.

the northern jet splits off into two branches at about 64°S, 0°E. While the northern branch flows in northwestward direction, the southern branch flows southwestward (Cisewski et al., 2011).

3.2.2. Moored current observations

The northernmost mooring, AWI-229, was positioned at the northwestern edge of the Maud Rise seamount at 64°S, 0°E and reveals the fluctuant current patterns of the Maud Rise Jet (MRJ). Daily mean horizontal current speeds measured at 64°S and vertically averaged between 100 and 200 m depth range between 0 and 24 cm s⁻¹ and show predominantly northwesterly to southwesterly flow (Fig. 5a). Mooring AWI-231 was placed southwest of Maud Rise where recirculation occurs; here the velocities range between 0 and 9 cm s⁻¹ and reveal northeasterly to southeasterly flow (Fig. 5b). Mooring AWI-232, which is located at the northern flank of the AntCC that follows the Antarctic continental slope. documents a strongly directional, northwestward flow running parallel to the depth contours (Figs. 5c and 6). Daily mean horizontal current speeds measured at 69°S, 0°E and spatially averaged between 100 and 200 m depth range between 0 and 13 cm s^{-1} and reveal a persistent northwesterly flow (Fig. 5c).

In order to identify oscillatory signals in the ADCP time series data we computed the power spectral density (PSD) of the horizontal and vertical velocities for each deployment period. In our analysis, we focused on the four major semi-diurnal and diurnal tidal constituents M₂ (12.42 h, 1.93 cpd), S₂ (12 h, 2.00 cpd), O₁ (25.82 h, 0.93 cpd) and K₁ (23.93 h, 1.00 cpd), which account for about 70-80% of the total tidal elevation in the Weddell Sea (Robertson et al., 1988). Moreover, rapid changes of wind at the sea surface often produce currents near the local inertial period, which ranges between 12.82 and 13.31 h at our mooring sites. At the two northernmost moorings (Fig. 7a and b), the power spectra of the horizontal current speeds reveal clear maxima of energy in the vicinity of the semidiurnal tides M₂ and S₂ and inertial periods. At the southernmost mooring (Fig. 7c), the layer-averaged power spectra show peaks related to both the semi-diurnal and diurnal tidal constituents. The PSDs of the vertical velocity (Fig. 8a-c) have a much flatter spectral slope - roughly one order of magnitude instead of 2.5 over the considered range of frequencies - than the horizontal speed PSDs, and show the highest peak at a frequency of about 1 cpd. This peak at 1 cpd, which dominates the spectrum of w, misses almost completely, or is at least insignificant, in the spectra of u and v. This suggests that the dominant diurnal cycle in vertical velocity is caused by something else than by tides. In the following chapters, we will demonstrate that this clear peak reflects the diel vertical migration of several zooplankton species. It is also interesting to note that the semidiurnal spectral peak of w is located exactly at S₂, i.e. is solar-cycle related,



Fig. 6. Detailed bottom topography of the Antarctic continental shelf slope between 1°W and 1°E, depicted from the Bathymetric Chart of the Weddell Sea (AWI BCWS), Antarctica (Schenke et al., 1998). Contours are every 200 m. The position of the southernmost mooring AWI-232 is indicated by the black dot and the 3-year mean current vector (vertically averaged between 100 and 200 m water depth) is shown by the black arrow. At 69°S the Antarctic Coastal Current features a strongly directional, northwestward mean flow of 6.2 cm s⁻¹ running parallel to the local depth contours.

while the semidiurnal spectral maxima of u and v are rather located at T and/or M₂.

3.3. Sea ice coverage and chlorophyll

Colour-coded mean sea ice concentration maps for the austral spring and summer months November until March, derived from SSM/I data applying the Bootstrap algorithm, are shown in Fig. 9. These maps reveal the seasonal decay of the sea ice cover and its interannual variability in the study area during the sampling period. The region is characterised by a recurring pattern of the vernal sea ice opening that begins with a polynya in vicinity of Maud Rise, which was first identified and described by Comiso and Gordon (1987). Between 2005 and 2007 the period of sea ice cover lasts 199 (54%), 223 (61%) and 275 (75%) days per year on average at 64°S, 66.5°S and 69°S (Table 4), respectively. The longest period of sea ice coverage was observed at all mooring locations in 2007. Based on a comprehensive ULS data set, Behrendt et al. (2011) showed that the mean sea ice draft along the Greenwich Meridian is relatively constant across the Weddell Gyre and only increases towards the coast. The authors observed similar mean sea ice drafts of 0.64 and 0.65 m and periods of ice cover of 241 and 246 days around Maud Rise at 64°S and 66.5°S. However, towards the Antarctic coast at 69°S the mean sea ice draft is much higher with 1.7 m for a period of 320 days. We compared the satellite-based sea ice concentrations derived from Bootstrap algorithm and ULS estimates of sea ice in the Lazarev Sea. Our results

show in agreement with Connolley (2005) that the Bootstrap data is largely in line with the ULS data, e.g. that the end of the sea ice season differs only by 1–6 days between both data sources.

Colour-coded¹ monthly mean maps of the chlorophyll a surface concentration, derived from satellite remote sensing and shown in Fig. 10, reveal the development and decay of phytoplankton blooms in the region during the sampling period. In the summer 2004/2005, a distinct algae bloom developed near Maud Rise with highest Chl concentrations around the mooring location at 66.5°S. The strongest $(Chl_{Max} > 4.5 \text{ mg m}^3)$ and largest bloom occurred in the summer 2005/2006, extending over all mooring sites for 3-7 weeks after the sea ice melt (Fig. 12). These findings were corroborated by in-situ measurements conducted during R.V. Polarstern cruise ANT-XXIII/2. The chlorophyll a concentration, which was measured at the end of the first mooring deployment along the prime meridian, reveals elevated phytoplankton abundance in the upper 40 m and a pronounced phytoplankton bloom of up to 4.2 mg m^{-3} chlorophyll between 65°S and 67°S (Fig. 11d). During winter 2006, the chlorophyll levels were below 0.1 mg m⁻³ along the entire transect (Fig. 11h). In the following summer 2006/2007, the observed chlorophyll concentrations were below 0.3 mg m^{-3} at all three mooring sites. Only a small bloom developed eastward of Maud Rise. In the summer 2007/2008, elevated chlorophyll levels at 64°S and 69°S indicate the development of a new ice edge bloom (Fig. 111).

3.4. Inference of diel vertical zooplankton migration from ADCP data

Mean 24-h cycles of the MVBS vertical distribution (with each cycle derived from 1 week of data) for the 155 consecutive weeks are displayed in Figs. 13, 15 and 17, which exhibit a variety of both distinct and diffuse bands of high backscatter. In order to illustrate and analyse the characteristic patterns in the temporal and vertical distribution of MVBS, we have selected twelve different weekly averaged diel patterns of the complete time series that are centred at seasonally distinctive dates: the autumnal equinox, the winter solstice, the vernal equinox and the summer solstice. These are shown together with the corresponding Doppler-derived vertical velocities, *w*, in Figs. 14, 16 and 18.

What stands out eyeing these overall 36 pairs of MVBS-w subfigures is a close match of the dominant patterns of both parameters. The highest downward vertical velocities always coincide with a downward sloping layer of enhanced backscatter and/or a clearance of the water column of scatterers starting at the top, and the highest upward vertical velocities go along with an upward sloping backscatter layer and/or a filling of the water column with scatterers from below. The conclusion from this observation is that the streaks of high vertical velocity are caused by the motion of assemblages of scatterers. Figs. 14, 16 and 18 also reveal a clearly diel cycle in the prominent patterns of MVBS and w. A distinct diel cycle in *w* was identified before by spectral analysis (Fig. 8). What the spectra however did not reveal is that the daily timing of upward or downward maximum vertical velocity is closely coupled to sunrise and sunset, hence changes with season (detailed later in Section 3.5). An exception is however the summer season, when a diel pattern is neither identifiable in the distribution of MVBS nor in w. Because there is no known physical process that could explain these patterns in backscatter and in w and their coupling to the daylight cycle, they are likely of biological nature. Since both the backscatter layers and the streaks of high w appear rather as contiguous signatures than as features, which can be related to discrete singular acoustic targets, we argue that they result from the bulk properties and behaviour of numerous smaller

¹ For interpretation of colour in Fig. 10, the reader is referred to the web version of this article.



Fig. 7. Power spectral densities (PSD) computed using horizontal current speeds from the moored ADCPs, which were vertically averaged between 100 and 200 m depth, for the deployments at 64°S (a), 66.5°S (b), and 69°S (c). Individual peaks at tidal constituent frequencies (O₁, K₁, M₂ and S₂) and local inertial frequency (T) are labelled.

animals within the ensonified vertical bins along the ADCP acoustic beams.

A more detailed look at Figs. 14a, e, i; 16a, e, i; and 18a, e, i, representing the autumnal equinox, allows to identify a nocturnal migration pattern of at least two distinct groups of migrators that is present at all three mooring sites and during all three sampled years 2005 through 2007. Generally, the one group, which we term deep migrators, is characterised in both MVBS and Doppler vertical

velocity data by quick migrations that are completed within roughly two hours during dawn and dusk. They leave the surface layer two hours before sunrise, descend to their daytime residence depth, and again attain the surface layer two hours after sunset. The daytime residence depth of the deep migrators during the autumnal equinox is revealed only in the ADCP record from 2006 at the southernmost mooring. Then and there it was found around 470 m depth (Fig. 18e). In all other records, the daytime residence



Fig. 8. Power spectral densities (PSD) computed using vertical velocities from the moored ADCPs, which were vertically averaged between 100 and 200 m depth, for the deployments at $64^{\circ}S$ (a), $66.5^{\circ}S$ (b), and $69^{\circ}S$ (c). Individual peaks at tidal constituent frequencies (O_1 , K_1 , M_2 and S_2) and local inertial frequency (T) are labelled.

depth of the deep migrators was below the maximum ADCP depth, even when the ADCP range extended down to 500 m. During nighttime the deep migrators were found occupying depths of 50 m or shallower. According to the method introduced by Cisewski et al. (2010) and in order to determine the descend and ascend speeds of the scattering layers, the inclined MVBS maxima associated with these migrators were fit by a hyperbolic tangent function. Mean and maximum downward migration velocities estimated this way from the slope of the curve are -3.0 cm s^{-1} and -8.0 cm s^{-1} . The steepest downward and upward trajectories of the deep migrators corresponded with the highest Doppler velocities, ranging between $\pm 2.0 \text{ cm s}^{-1}$ and $\pm 3.0 \text{ cm s}^{-1}$. The second group consists of "slow" migrators descending from the surface layer to a depth ranging between 250 m and 350 m one hour later. This group reaches its residence depth at noon and attains the uppermost 50 m one hour after sunset. The mean and maximum



Fig. 9. Colour-coded sea ice concentration maps from SSM/I data using the Bootstrap algorithm reveal monthly mean ice concentration within the study area for the months January, February, March, November and December of the consecutive years 2005–2008. These maps show the seasonal and interannual variability of the sea ice concentration i.e. exhibiting the decay and the new development of sea ice near Maud Rise during the sampling period. Bootstrap algorithm data (Comiso, 1999) were retrieved from NSIDC via http://www.nsidc.org/data/nsidc-0079.html. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Table 4

Periods of sea ice coverage at the mooring sites derived from ULS and SSM/I bootstrap data.

Latitude Year	Ice coverage from ULS	Mean Ice coverage from SSM/			Mean
	Period	Days	Period	Days	Days
2005	07/06/2005-12/16/2005	161	06/08/2005-12/19/2005	195	
2006	n.a.	n.a.	05/27/2006-12/05/2006	193	199
2007	n.a.	n.a.	05/28/2007-12/21/2007	208	
2005	05/31/2005-12/15/2005	199	05/23/2005-12/09/2005	201	
2006	05/24/2006-12/13/2006	204	05/10/2006-12/10/2006	215	223
2007	05/16/2007-01/08/2008	238	05/03/2007-01/10/2008	254	
2005	n.a.	n.a.	05/02/2005-12/20/2005	233	
2006	04/14/2006-01/20/2007	282	04/06/2006-01/21/2007	291	275
2007	04/14/2007-01/19/2008	281	03/31/2007-01/23/2008	300	
	Year 2005 2006 2007 2005 2006 2007 2005 2006 2007	Year Ice coverage from ULS Period 2005 07/06/2005-12/16/2005 2006 n.a. 2007 n.a. 2006 05/31/2005-12/15/2005 2006 05/24/2006-12/13/2006 2007 05/16/2007-01/08/2008 2005 n.a. 2005 n.a. 2006 05/24/2006-12/13/2006 2007 05/16/2007-01/08/2008 2005 n.a. 2006 04/14/2006-01/20/2007 2007 04/14/2007-01/19/2008	Year Ice coverage from ULS Mean Period Days 2005 07/06/2005-12/16/2005 161 2006 n.a. n.a. 2007 n.a. n.a. 2005 05/31/2005-12/15/2005 199 2006 05/24/2006-12/13/2006 204 2007 05/16/2007-01/08/2008 238 2005 n.a. n.a. 2005 n.a. n.a. 2007 05/16/2007-01/08/2008 238 2005 n.a. n.a. 2006 04/14/2006-01/20/2007 282 2007 04/14/2007-01/19/2008 281	Year Ice coverage from ULS Mean Ice coverage from SSM/I BT Period Days Period 2005 07/06/2005-12/16/2005 161 06/08/2005-12/19/2005 2006 n.a. n.a. 05/27/2006-12/05/2006 2007 n.a. 05/28/2007-12/21/2007 2005 05/31/2005-12/15/2005 199 05/23/2005-12/09/2005 2006 05/24/2006-12/13/2006 204 05/10/2006-12/10/2006 2007 05/16/2007-01/08/2008 238 05/03/2007-01/10/2008 2005 n.a. n.a. 05/02/2005-12/20/2005 2005 n.a. n.a. 05/03/2007-01/10/2008 2005 n.a. n.a. 05/02/2005-12/20/2005 2006 04/14/2006-01/20/2007 282 04/06/2006-01/21/2007 2007 04/14/2007-01/19/2008 281 03/31/2007-01/23/2008	Year Ice coverage from ULS Mean Ice coverage from SSM/I BT Period Days Period Days 2005 07/06/2005-12/16/2005 161 06/08/2005-12/19/2005 195 2006 n.a. n.a. 05/27/2006-12/05/2006 193 2007 n.a. 05/28/2007-12/21/2007 208 2005 05/31/2005-12/15/2005 199 05/23/2005-12/09/2005 201 2006 05/24/2006-12/13/2006 204 05/10/2006-12/10/2006 215 2007 05/16/2007-01/08/2008 238 05/03/2007-01/10/2008 254 2005 n.a. n.a. 05/02/2005-12/20/2005 233 2006 04/14/2006-01/20/2007 282 04/06/2006-01/21/2007 291 2007 04/14/2007-01/19/2008 281 03/31/2007-01/23/2008 300

downward/upward migration velocities estimated from the slope of the fitted parabolic function are ± 1.0 cm s⁻¹ and ± 2.0 cm s⁻¹, respectively.

The nocturnal migration patterns of the deep migrators are again apparent in the vertical distribution of MVBS and vertical velocity at all three mooring sites around the winter solstice. At 64°S, the DVM pattern is characterised by clearance of the upper water column (e.g. Fig. 14b and j). In 2005, clearance starts three hours before sunrise and extends to more than 300 m three hours later around noon (\sim 11:30 GMT). Ascent occurs during the following 4.5 h. At 66.5°S, the deep migrators occupy the layer between 75 and 125 m water depth at night, yielding a strong scattering layer (Fig. 16b, f and j). They leave the surface layer three hours before sunrise, descend quickly to a daytime residence below the ADCP depth, and attain their nighttime residence layer again three hours after sunset. Further south at 69°S, the MVBS distribution



Fig. 10. Monthly mean surface concentration of chlorophyll a within the study area for the months January, February, March, November and December of the consecutive years 2005–2008. These maps were acquired using products processed and distributed by ACRI-ST GlobColour service, supported by EU FP7 MyOcean & ESA GlobColour Projects, using ESA ENVISAT MERIS data, NASA MODIS and SeaWiFS data.

reveals one distinct group of deep migrators, which occupy the layer between 170 and 200 m depth during the night (Fig. 18b, f and j). Although the sun stays under the horizon throughout the day around winter solstice, the deep migrators descend to a depth of around 450 m starting at 9:00 GMT (Fig. 18f and j). Mean and maximum slope velocities of the corresponding scattering layer are $\pm 2.0 \text{ cm s}^{-1}$ and $\pm 3.0 \text{ cm s}^{-1}$. After reaching their residence depth around noon, they start to ascend soon after and attain their shallow nighttime residence depth again at 15:00 GMT.

Nocturnal diel migration is also apparent in the MVBS and vertical velocity data at all three mooring sites around the vernal equinox. Since the earth experiences equal hours of daylight and darkness during autumnal and vernal equinox, the observed patterns and timing of DVM are very similar during these periods. For example at 66.5°S, the deep migrators reveal a similar nocturnal DVM pattern around the autumnal and the vernal equinox. In both seasons of the years 2005 and 2006, they leave the surface layer two hours before sunrise, descend quickly to a daytime residence below the ADCP depth, and attain the surface layer two hours after sunset (Fig. 16a, c, e and g). DVM around the vernal equinox however is consistently weaker in 2007 than in the two other years at all three mooring locations.

While typical nocturnal migration patterns of two distinct groups of migrators are recognisable in both backscatter and Doppler-shift vertical velocity data observed around autumnal equinox, winter solstice and vernal equinox, the DVM ceased between mid-November and the summer solstice within the observed depth range at all three mooring sites (Figs. 14d, h, l; 16d, h, l and 18d, h, l). However, no statement can be made about what happened in the top few tens of metres. In all years, the summer distribution of MVBS reveals an approximately vertically layered structure with constant depths of the scattering layers and without indications of DVM. During summer solstice 2005/2006 at 66.5°S and 69°S (Figs. 16d and 18d), highest MVBS values are found above 50 m depth and lowest values below 300 m. During the other summers and at 64°S, the scatterers appear distributed more evenly over the water column or concentrated at several deeper depth levels.

3.5. Seasonal and interannual variation of the vertical velocity

In order to analyse the seasonal and interannual variability of the migration velocity, mean diel cycles of the Doppler-derived vertical velocity (averaged between 100 and 200 m depth) are calculated for successive months between February 2005 and February 2008. This analysis also considers the times of local sunrise and sunset at the different mooring positions to examine the role of the astronomical daylight cycle. Fig. 19a–d document that the temporal distribution of the ascent and descent velocities at 64°S peak symmetrically around noon and reveal a clear dependence on the sun angle between January/February and October/November of the years 2005–2007 at the northernmost



Fig. 11. Summer 2005/2006 sections (a) of temperature, (b) salinity, (c) potential density and (d) chlorophyll. The data were collected during R.V. *Polarstern* cruise ANT-XXIII/ 2 (December 14 and December 20, 2005). Winter 2006 sections (e) of temperature, (f) salinity, (g) potential density and (h) chlorophyll. The data were collected during R.V. *Polarstern* cruise ANT-XXIII/6 (July 11 and August 11, 2006). Summer 2007/2008 sections (i) of temperature, (j) salinity, (k) potential density and (l) chlorophyll. The data were collected during R.V. *Polarstern* cruise ANT-XXIII/6 (July 11 and August 11, 2006). Summer 2007/2008 sections (i) of temperature, (j) salinity, (k) potential density and (l) chlorophyll. The data were collected during R.V. *Polarstern* cruise ANT-XXIV/2 (January 17 and January 23, 2008). Moorings positions are illustrated as white and black dashed curves.

mooring. In parallel with the shortening of the day length between February and the winter solstice (June 21), the downward and upward migration peaks shifted by around ±1 h per month. The highest ascent and descent velocities of 1.6 cm s⁻¹ and -1.3 cm s⁻¹ were observed in March 2005. A pronounced change of the vertical velocity pattern occurs from mid-November 2005 to January 2006. During this period, the synchronised diel vertical migration ceases. In February 2006 (Fig. 19b), the nocturnal DVM resumes and lasts to November 2006. However, the monthly average maximum ascent and descent velocities of about 1.0 cm s⁻¹ and -0.5 cm s⁻¹ were smaller than those observed 2005. Between December 2006 and January 2007, no DVM patterns are apparent. In January 2007 (Fig. 19c), DVM resumes and the highest ascent and descent velocities of 1.4 cm s^{-1} and -1.4 cm s^{-1} , occurring between March and May, are comparable to those observed in 2005. Between October 2007 and January 2008 DVM ceases again and starts once more in February 2008 (Fig. 19d).

The migration velocities observed at 66.5°S are very similar to those observed at 64°S, with peaks distributed symmetrically around noon, revealing a clear dependence on the sun angle between February and October of the years 2005–2007 (Fig. 20a–d). The highest ascent and descent velocities of 2.1 cm s⁻¹ and -1.8 cm s⁻¹ were observed in March/April 2005. Between July and October, the maximum ascent and descent velocities range

between 1.1 cm s^{-1} and 0.6 cm s^{-1} , and -1.0 cm s^{-1} and -0.6 cm s^{-1} . In agreement with the observation made at 64°S, a change of the vertical velocity pattern occurs from mid-November 2005 to January 2006 when the diel vertical migration ceases. In February 2006 (Fig. 20b), the nocturnal DVM starts again and lasts until November 2006. However, the peak ascent and descent velocities of -0.8 cm s^{-1} and 0.9 cm s^{-1} were smaller than those observed 2005. Between October 2006 and January 2007, no DVM patterns are apparent. In February 2007 (Fig. 20c), DVM resumes and the highest ascent and descent velocities of 1.6 cm s^{-1} are comparable to those observed the same month in 2005. Between November 2007 and January 2008 DVM ceases and resumes in February 2008 (Fig. 20d).

The location of the southernmost mooring at 69°S is exposed to extreme variations in daylight hours, with twenty-four hours of daylight during summer (November 23 to January 19) and the sun below the horizon around mid-winter for the period May 31 to July 13. The ascent and descent velocities are distributed symmetrically around noon and reveal a clear dependence on the sun angle between February and April/May of the years 2005–2007 (Fig. 21a–d). However, DVM continues through the polar night until October in all three years. With the sun under the horizon, descent occurs around 8 a.m. and ascent around 3 p.m. local time. The maximum ascent and descent velocities in the course of the



Fig. 12. Time series of chlorophyll a concentration estimated from 8-day composites of merged MERIS/MODIS/SeaWiFS data (upper panel) and ULS derived ice coverage (black line) compared to SSM/I derived ice coverage (red line) (lower panel) at 64°S (a), 66.5°S (b) and 69°S (c). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

year of 1.5 cm s^{-1} and -1.2 cm s^{-1} were observed between February and April of the years 2005–2007. During the entire observation period (February 2005 to March 2008), DVM ceases between November and January and resumes in February of the following year.

The basic DVM pattern, which evolves from synopsis of the Doppler-*w* measurements taken through the whole three-year long deployment periods at all three mooring locations, can be summarised as follows: maximum downward and upward vertical migration speeds are distributed symmetrically around local noon; downward migration occurs during the two hours before sunrise and upward migration during the two hours after sunset; when the sun does not rise above the horizon during the polar night descent occurs around 8 a.m. and ascent around 3 p.m. local time. The annual modulation of the DVM in contrast is asymmetric relative

to the summer and winter solstices; vertical migration speeds are highest during late summer and early autumn (February until April); DVM persists throughout winter, but the vertical speeds tend to decline between May and October; in late spring/early summer DVM ceases and is then suspended until high summer, for the months November through January.

3.6. Seasonal and interannual variation of the integrated mean volume backscattering strength

A different aspect of the observed asymmetry over the annual cycle is revealed by considering temporal changes in the mean zooplankton concentration in the water column, deduced from vertical means of normalised MVBS profiles. In order to investigate the occurrence and the seasonality of relative changes of zooplankton



Fig. 13. Annual time series of weekly averaged 24 h cycles of mean volume backscatter strength at 64°S in the 155-week data series from 10 February 2005 to 21 February 2008 ("midnight" is signified by tick marks on the horizontal axis).

backscatter, we applied a normalisation to the acoustic data. To examine the annual cycle, daily means of MVBS, averaged between 50 and 300 m water depth, were calculated for all mooring sites. We define $MVBS_{med}$ as the median value (in the linear domain) of all available daily averages of MVBS values averaged between 50 and 300 m water depth. The difference between MVBS and MVBS_{med} in the logarithmic domain gives the normalised MVBS value: $MVBS_{norm} = MVBS - MVBS_{med}$. Generally, the time series of the normalised MVBS declined at all sites (64°S, 66.5°S and 69°S) during autumn towards winter and increased throughout the spring with greatest levels observed during summer/early autumn (Fig. 22). However, the magnitude of the observed decline/increase in $MVBS_{50-300m}$ was different between the locations and varied from year to year. Fig. 22 shows that the largest autumnal decline occurred at 64°S, where the difference between the summer maximum (2005/2006) and the winter minimum (2006) amounted to nearly 14 dB. The three-year maximum at 64°S of about 7 dB was observed in February 2006, followed by a 14 month lasting decreasing trend to about -5 dB in May 2007 and a re-increase during autumn towards winter to a maximum of about 4 dB in January/February 2008. The normalised MVBS declined at 66.5 and 69°S during autumn towards winter/early spring and re-increased throughout the spring with greatest levels observed during summer/early autumn.

Averaging the daily normalised MVBS values obtained from the three years of measurements yields a representation of the mean annual cycle of relative changes of zooplankton backscatter (Fig. 23). Accordingly, the highest value at 64°S occurs in February, at 66.5°S in December and at 69°S in April, on average over all three latitudes thus during the season summer to early autumn. The lowest value at 64°S is found in June, at 66.5°S in August and at 69°S in October, on average over all three latitudes thus during the season winter to spring. Fig. 23 however also reveals that the annual averages are associated with a large standard deviation, which is at least partly related to interannual variability. In case of the mooring position 64°S, the differences between the three years are so large that the overall deployment length apparently is still not long enough to obtain what can be considered a regular mean annual cycle.

4. Discussion

4.1. Seasonal and interannual variability of DVM

Our 3-year long moored ADCP time series of MVBS and vertical velocity have demonstrated diel vertical migration (DVM) to be a persistent behavioural pattern at all three mooring sites that is



Fig. 14. Mean diel cycle of the sun angle (top row), mean volume back scattering strength MVBS (2nd, 4th and 6th rows from top) and Doppler vertical velocity w (3rd, 5th and 7th row from top) at 64°S, 0°E estimated for four different weeks near autumnal equinox, winter solstice, vernal equinox and summer solstice of the years 2005, 2006 and 2007.

closely related to the astronomical daylight cycles. The vertical migration of zooplankton however changed in late spring/early austral summer at all three locations. In contrast to many previous acoustic studies in other ocean regions, DVM at our observation sites continued throughout winter, even at the highest latitude exposed to the polar night, until late spring (late October/early November) when it ceased completely.

Examination of the temporal change of scattering layers revealed two distinct groups of migrators at all three mooring positions: (i) deep/fast migrators: species migrating from near the surface to below the ADCP depth (>400 m) at dawn and from their daytime residence depth to the surface at dusk and (ii) shallow/ slow migrators: species migrating from near the surface to 350 m at dawn and to the surface at dusk (Cisewski et al., 2010). Vertical migration speeds were estimated from the slopes of the scattering layers and from the vertical velocity components of the ADCP, yielding average speeds of $\pm 2.0-3.0$ and $\pm 1.0-2.0$ cm s⁻¹ and maximum speeds of ± 8.0 cm s⁻¹ and ± 3.0 cm s⁻¹, respectively. Thus, our analysis has shown that the mean and maximum migration speeds estimated by the vertical component of the ADCP velocity were generally lower than those estimated from the slopes of the different sound scattering layers. This is consistent with other ADCP observations in the North Pacific, Northeast Atlantic, Arabian Sea and Ligurian Sea, reported by Plueddemann and Pinkel (1989), Heywood (1996), Luo et al. (2000) and Tarling et al. (2001), respectively. While the migrations speeds estimated by fitting a curve to the undulating sound scattering layers represent an estimate of the speed of the fastest coherently migrating scatterers alone, the vertical velocities from the ADCP measure a slightly different mean speed that comes from an intensity-weighted sum of the scatterer velocities within the ensonified volume (Plueddemann and Pinkel, 1989). The same authors concluded that the vertical ADCP velocities can be expected to underestimate the true migration rates in regions where the ratio of the intensity of migrating scatterers to that of non-migrating scatterers is small.

While our acoustic study of the seasonal occurrence, daily timing, and pattern of diel vertical migration, based on three years of acoustic measurements from three locations, is - to our knowledge - unique for the Weddell Sea, some similar studies have been conducted in other ocean regions. From the Southern Ocean too is a study based on a multitude of Drake Passage transits by a ship equipped with a 150 kHz ADCP (Chereskin and Tarling, 2007), which also identified DVM as a dominant source of variability, and that DVM continued through winter. However, these Drake Passage transits did not extend to south of the polar circle. In the central Greenland Sea, moored ADCPs were deployed for one year and exhibited a diurnal cycle typical for vertically migrating zooplankton both in vertical velocity and in acoustic backscatter (Fischer and Visbeck, 1993). Peak ascent and descent velocities were around ± 1.5 cm s⁻¹. Strong seasonal variations in the DVM were evident, and the timing as well as the migration amplitude changed with daylight as the season progressed. In summer and



Fig. 15. Annual time series of weekly averaged 24 h cycles of mean volume backscatter strength at 66,5°S in the 158-week data series from 10 February 2005 to 28 February 2008 ("midnight" is signified by tick marks on the horizontal axis).

during the polar night the migration became very weak and was only detectable in the displacement of scattering layers. Benoit et al. (2010) used continuous multifrequency echosounding validated by trammel net captures in the Arctic Ocean to analyse the vertical migration of polar cod in relation to the photoperiod, the thermal structure of the water column and the vertical distribution of its main calanoid copepod prey. The DVM stopped in May coincident with the midnight sun and increased schooling and feeding. Wallace et al. (2010) analysed a two-year time series of acoustic backscatter and velocity data from moored ADCPs obtained in an ice-free and a seasonally ice-covered Arctic fjord. They observed a strong seasonal cycle in migratory behaviour in both fjords, with classic DVM apparent in spring and autumn. However, differences in vertical migration behaviours emerged during summertime, with no synchronous signal observed under ice-free conditions, whereas the seasonally ice-covered fiord exhibited both asynchronous and weakly synchronous migration patterns.

4.2. Environmental factors affecting DVM

4.2.1. Light

4.2.1.1. DVM during polar night. Generally assumed is that DVM at high latitudes slows down or ceases during wintertime as food

availability decreases and some zooplankton species enter diapause and so spend the winter in the deep, dark and presumably safer interior of the ocean (Atkinson, 1998; Ashjian et al., 2003). Our data but show that DVM persisted throughout winter even at the southernmost mooring with only somewhat reduced vertical migration speeds. A similar observation was made by Berge et al. (2009), who presented acoustic data from two coastal locations in Svalbard (Kongsfjorden and Rijpfjorden at 79°N and 80°N, respectively) that indicate continuation of synchronised DVM of zooplankton throughout Arctic winter, in both open and icecovered waters. The authors argue that even during the polar night DVM is regulated by diel variations in solar and lunar illuminations, which are at intensities far below the threshold of human perception. The observed DVM signal was greater under ice-free conditions than under ice-covered conditions, possibly explicable through attenuation of light by the sea-ice cover that weakens the intensity of $\Delta I/I$ in the under-ice water column.

Our data challenge the view that daylight exerts a strong exogenous cue when the sun stays under the horizon throughout the day around winter solstice and the surface is covered by sea ice. Our MVBS records reveal one distinct group of deep migrators at 69°S, which occupy the layer between 170 and 200 m depth during the night and descend to below 450 m during the day, thus a DVM



Fig. 16. Same as Fig. 14 but for the mooring located 66.5°S.

between much deeper and hence darker layers than the mere top 100 m investigated by Berge et al. (2009). The main difference between winter and the other seasons at the polar latitude in our observations is a deeper night-time sojourn level, 170–200 m in winter instead of 100 m or shallower from spring through autumn. This observation suggests that the deep migrators in winter are more attracted by a pelagic food source than by a food source that is associated with the sea ice. A possible explanation for this behavioural change is a gradual move from herbivorous nutrition during spring through autumn to carnivorous foraging during winter, assuming that the smaller and less mobile prey organisms are more closely tied the upper part of the water column, hence the mixed layer in which the phytoplankton occurred during summer (Flores et al., 2014).

The deep DVM during the polar night in our results suggests a biological clock as Zeitgeber. In a recent study, Teschke et al. (2011) showed that *Euphausia superba* possesses an endogenous circadian clock that governs metabolic and physiological output rhythms. Kawaguchi et al. (1986) and Meyer et al. (2010) demonstrated that one of the overwintering strategies for adult krill is to reduce feeding and metabolic activity during winter. However, the decline of metabolic activity is hypothesised to be not only a response to reduced food availability but also driven by an endogenous timing system synchronised by the photoperiod (Teschke et al., 2011; Meyer, 2012). An existence of biological clocks is also suggested by observations made in the deep North Atlantic below 500 m depth, showing tuning of DVM to latitudinal and seasonal

changes in day length despite light being an implausible cue (van Haren and Compton, 2013).

4.2.1.2. DVM during midnight sun. While DVM persists from February to October, the zooplankton communities cease their migration in the investigated depth range beginning late October/early November. During this transition period, the light environment is changing from a true day-night contrast to one of continuous sunlight, depending on latitude either dim during the night or with the sun well above the horizon throughout. The observed halt of DVM around October/November could have been caused either by fewer animals choosing to migrate or by a decrease in animal abundance. However, the time courses of mean normalised mean volume backscattering strength in the upper 50-300 m (Fig. 22) do not indicate a particular decrease of animal abundance at this time of the year. In the three-year mean, integrated backscattering strength generally declined at all sites (Fig. 23) during autumn towards winter and increased throughout spring and summer. Because the halt of DVM in late spring/early summer thus cannot be explained by a sudden lack of animals, it must therefore be related to a change in their behaviour. In a recent study, Flores et al. (2014) investigated the macrozooplankton and micronekton community of the Lazarev Sea at three depth layers (0-2 m; 0-200 m; and 0-3000 m) during austral summer, autumn and winter. The authors showed that abundant krill species like Thysanoessa macrura may not cease their DVM during summer but decrease the amplitude in shallower depths <50 m. However, such changes



Fig. 17. Annual time series of weekly averaged 24 h cycles of mean volume backscatter strenght at 69°S in the 156-week data series from 24 February 2005 to 28 February 2008 ("midnight" is signified by tick marks on the horizontal axis).

in the upper layer of the water column could not be detected by our moored ADCPs, because they did not sample above 20 m water depth.

The halt of DVM during summer was also reported for the Arctic region by Bogorov (1946), Buchanan and Haney (1980) and Blachowiak-Samolyk et al. (2006). Based on net tows, these authors showed that zooplankton populations did not migrate in ice-free waters under continuous light conditions in midsummer, because without the light/dark cycle endogenous rhythms in vertical movements cannot be entrained (Cohen and Forward, 2009). The absence of DVM has been attributed to a weak change in irradiance (Buchanan and Haney) and and/or vanishing diel variation of the threat from visually oriented predators during the quasicontinuous illumination during the polar summer (Hays, 1995). This is consistent with our results for the early summer period, November-mid December, when DVM did not occur and the observed zooplankton communities remained in the uppermost 50 m at all three sites, but not so for the late summer month February, when the astronomical light cycle was almost the same as nearly November and high vertical migration speeds were recorded.

In a recent study, Cottier et al. (2006) measured the DVM of zooplankton in an Arctic Fjord at 79°N between June and September 2002. During this period, the light environment changed from one

of continuous sunlight to a true day-night contrast. The vertical velocity was measured by a 300 kHz ADCP and indicated unsynchronized migration of individual animals during the weeks of continuous daylight and changing to synchronised DVM when true night-time cycles resumed toward autumn. Wallace et al. (2013) pointed out that a lack of synchronised DVM at high latitudes does not necessarily mean that DVM does not take place. Although zooplankton populations do not migrate vertically in a synchronised manner during midnight sun conditions, Cottier et al. (2006) and Wallace et al. (2010) found evidence that individuals within those populations performed forays in and out of the surface layers throughout the 24-h cycle. While their backscatter data indicated that there was no net vertical displacement of the population at any time during the 24-h period, the vertical velocity showed a continuous net downward movement in the surface layers and a net upward movement at depth, which represents the so-called unsynchronised DVM. In our study, we also analysed both ADCP backscatter and vertical velocity data, but we found no evidence for unsynchronised DVM during Austral summer, when the synchronised DVM ceases within the depth range.

4.2.2. Sea ice and phytoplankton blooms

Sea ice contributes to variability in high-latitude primary production by affecting light availability, ocean stratification and



Fig. 18. Same as Fig. 14 but for the mooring located 69°S.

nutrient dynamics; by serving as a substrate for concentrated algal biomass and growth; by generating phytoplankton blooms upon its melt during spring–summer with influence on whole ecosystems (Massom and Stammerjohn, 2010; Taylor et al., 2013); and by creating the sole habitat of certain species at the lowest and highest trophic levels. Figs. 9 and 10 illustrate the relationship between ice retreat and the spring bloom in the study area and show that the timing of the spring bloom and its extent vary from year to year, but without a conspicuous relation to the interannual variation of the vernal sea ice melt.

The strongest and largest bloom (Chl_{Max} > 4 mg m⁻³) occurred in the summer 2005/2006 and was observed at all mooring sites 3-7 weeks after the sea ice decayed (Fig. 12). During the same period, the distribution of the MVBS reveals an approximately vertically layered structure with no indications of DVM. Highest backscatter values were found at the uppermost layer at 66.5°S and 69°S (Figs. 16d and 18d). At 69°S, the sun does not set down between November 23 and January 19 of the following year. During the summer period, there is no obvious optimal time for zooplankton to visit the surface layers because the continuous illumination maintains the threat from visually oriented predation during day and night. Since the relative rate of change of illumination is minimal at this time, the ultimate causes of migration may become separated from the light-mediated proximal cue that coordinates it (Cottier et al., 2006). Thus, the increased food supply associated with the phytoplankton bloom may cause the zooplankton communities to cease their DVM in favour of feeding.

In summer 2006/2007, no marked phytoplankton bloom developed after the vernal sea ice melt in the Lazarev Sea. Satellite imagery only reveals a patch of increased surface Chl close to and east of the mooring location at 66.5°S. While DVM ceased as in the year before at all three mooring locations, the zooplankton according to the vertical MVBS distribution appeared staying deeper in the water column throughout the day than in the preceding summer (Figs. 14, 16 and 18). During the summer survey 2007/2008, the seasonal sea ice cover in December was less reduced than in the previous two summers while the chlorophyll distribution revealed a large bow-shaped phytoplankton bloom of $Chl_{Max} > 2 \text{ mg m}^{-3}$ bend around the northwestern edge of Maud Rise (Figs. 9 and 10). Interesting to note is that not only the moored ADCP time series document a ceasing of the DVM as in the two years before, but that also the data collected with the ship-mounted SIMRAD EK60 zooplankton echosounder collected during the survey did not show signs of a synchronised diel vertical migration (Brandt et al., 2011).

The 3-year mean annual time series of normalised MVBS_{50-300m} (Fig. 23), while still subject to irregularities that mainly result from interannual variability, reveal maximum values during summer and early autumn (December through April) and minima in late winter and spring (June through October). Such annual cycle can be interpreted as following the canonical change of primary production in the course of the year, exhibiting a phytoplankton bloom in spring/summer that creates the dominant annual food basis. According to the satellite maps of surface Chl (Fig. 10), the spring bloom peaks in January in our study area during the three



Fig. 19. Seasonal variation of the diel vertical migration velocity at 64° S, averaged between 100 and 200 m depth for the years 2005 (a), 2006 (b), 2007 (c), and 2008 (d). Contour intervals are 0.1 cm s⁻¹. Dashed lines show the times of local sunrise and sunset.

years observation period but differs greatly between years in terms of its duration and areal extent. Comparison of the normalised MVBS_{50–300m} time series from individual years also reveals considerable interannual variability (Fig. 22). The most pronounced annual maximum in all three year occurs in the summer

2005/2006, which is the summer at which also the strongest and largest phytoplankton bloom, affecting all the mooring locations, was indicated by satellite imagery. In summer 2006/2007 in contrast, a pronounced although short maximum of MVBS_{50-300m} is only indicated by the time series from 66.5°S, which is the only



Fig. 20. Seasonal variation of the diel vertical migration velocity at 66.5°S, averaged between 100 and 200 m depth for the years 2005 (a), 2006 (b), 2007 (c), and 2008 (d). Contour intervals are 0.1 cm s⁻¹. Dashed lines show the times of local sunrise and sunset.

location at which in this year a spatially confined phytoplankton bloom is indicated in satellite Chl maps (Fig. 10); the summer maximum of MVBS_{50–300m} at 69°S is only modest while it misses almost completely at 64°S in 2006/2007. At the end of our observations in

the summer 2007/2008 MVBS_{50-300m} increases at all latitudes but strongest at 64°S (Figs. 22 and 23), which is the only location affected by the bow-shaped phytoplankton bloom that occurred around the northwestern edge of Maud Rise that year. At the two



Fig. 21. Seasonal variation of the diel vertical migration velocity at 69°S, averaged between 100 and 200 m depth for the years 2005 (a), 2006 (b), 2007 (c), and 2008 (d). Contour intervals are 0.1 cm s⁻¹. Dashed lines show the times of local sunrise and sunset.

more southern mooring locations the bloom appears delayed until March, 2008 (Fig. 10). Together, comparison of the individual time series of MVBS_{50-300m} from the three mooring locations and of the surface Chl distribution in the area reveals a positive correlation between zooplankton biomass implied by MVBS_{50-300m} and

phytoplankton biomass indicated by Chl. The question is what controls phytoplankton biomass?

During both summer surveys, the impact of the sea ice on the initiation of the spring bloom in the Lazarev Sea was examined using discrete hydrographic, sea ice and chlorophyll data. The



Fig. 22. Time series of normalised MVBS in the depth range 50–300 m (daily averages as red line and 7-day running mean as black line): (a) at 64°S, (b) at 66.5°S and (c) at 69°S. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

melting of the sea ice along the Greenwich Meridian in summer 2007/2008 is illustrated by warming and freshening of the uppermost 50 m of the Antarctic Surface Water layer (Fig. 11i–k). After the sea ice retreat, low-salinity melt water stabilizes a shallow mixed-layer that is generally favourable for phytoplankton primary production. However, while the observed phytoplankton bloom (Fig. 11l) occurred in shallow mixed layers down to 40 m depth between 64° and 62°S left after the sea ice melt, not all melt water lenses supported phytoplankton blooms.

In summary, neither the sea ice coverage at the end of winter nor its vernal retreat and the associated mixed layer shallowing can hence be considered the primary determinator of the timing and magnitude of the seasonal phytoplankton bloom. Another bottom up control on phytoplankton primary production could possibly be exerted by the availability of the trace nutrient iron. Iron plays an essential role in photosynthesis and has been demonstrated by in-situ experiments to strongly influence phytoplankton growth in the Southern Ocean (e.g. Smetacek et al., 2012). Due to lack of data on iron we are unfortunately unable to investigate the impact of its availability on our observations. However, as an alternative to bottom up control we hypothesise top down control by zooplankton grazing on phytoplankton as a partial explanation for the observed interannual variations in the phytoplankton blooms magnitude. The highest values of MVBS_{50-300m} at all three mooring locations occurred in late summer 2005/2006 (Fig. 22), following the big phytoplankton bloom in that season (Fig. 10). During the subsequent autumn and winter, MVBS_{50-300m} gradually declined. But originating from the high biomass accumulated in summer 2005/2006, higher MVBS_{50-300m} values remained until the next spring, 2006/2007, than were recorded the spring before.

In order to survive the dark season when photosynthetic primary production is insufficient to balance the metabolic losses of heterotrophs, polar zooplankton have developed several overwintering strategies. These strategies, comprising inter alia opportunistic feeding on varying sources, combustion of body substance and degrowth, and dormancy or diapause, can differ between groups, species and even developmental stages (Clarke and Peck, 1991; Bathmann et al., 1993; Torres et al., 1994; Hagen and Schnack-Schiel, 1996; Meyer, 2012; Auerswald et al., 2015). Many crustaceans for instance, in particular the polar species, are able to store energy in the form of lipid reserves. Copepods can build up massive amounts of lipids exceeding 50% of their dry mass, some of the highest lipid levels in organisms on earth (Kattner and Hagen, 2009). For Antarctic krill, *Euphausia superba*,



Fig. 23. Annual mean time series of normalised MVBS (annual averages ± standard deviation) in the depth range 50-300 m (a) at 64°S, (b) at 66.5°S and (c) at 69°S.

lipid production also appears effective enough to accumulate large energy reserves for winter. Changes in lipid content of krill over the phytoplankton growth season were found most pronounced in the immature and adult specimens, increasing from about 10% lipid of dry mass in late winter/early spring to more than 40% in autumn (Hagen et al., 2001). Regarding the consumption of body energy reserves of krill during one particular winter in the Lazarev Sea, Schmidt et al. (2014) report a reduction of the lipid content of the digestive gland from 45% of dry mass in April, 2004, to 8% in December, 2005.

Zooplankton surviving the winter 2005–2006, hence abundant in spring 2006/2007, thus may have grazed down the phytoplankton when it started to grow, preventing it from forming a substantial bloom. In turn, due to curbing the phytoplankton bloom zooplankton found less food than the preceding summer and was not able to build up so much biomass than the summer before. With zooplankton being consequently less abundant the following spring, a moderate phytoplankton bloom could again develop in summer 2007/2008.

4.2.3. Hydrography and circulation

While differences between the mooring sites are evident if the records of mean backscatter are compared (Fig. 22), the time series on the whole are dominated by very similar patterns of diel displacements of the MVBS signatures and of the vertical migration velocities (Figs. 14, 16 and 18). A likely explanation for the observed similarity is the regional hydrographic regime. All three moorings are located within the same large-scale circulation system, the southern limb of the Weddell Gyre in the Lazarev Sea, with the transport of water masses governed by two comparably narrow current branches, the MRJ bound to the lower continental slope directly north of Maud Rise and the AntCC confined to the Antarctic continental slope (Cisewski et al., 2011). The flow field between these two current cores is, in its northern part, influenced by the stagnant Taylor column formed on top of Maud Rise

and elsewhere rather sluggish, partly recirculating eastward, and dominated by transient mesoscale eddies. The sluggish flow probably explains that the spatially limited phytoplankton bloom occurring during summer 2006/2007 around 66.5°S is reflected by a MVBS_{50-300m} peak (Figs. 10 and 22) at this location. We are however not arguing that this MVBS_{50-300m} increase is solely due to zooplankton growth; rather invasion of zooplankton by use of vertical shears in the flow field and adaptation of the DVM migration depth to food in the mixed layer, as described in Krägefsky et al. (2009) may have made the major contribution. Leach et al. (2011) demonstrated that the eddies, mostly shed by instability of the MRJ, and their associated either warm or salty cores, gradually mix. The hydrographic regime between the moorings, although spread over a meridional distance of more than 550 km, is dominated by the inflowing Warm Deep Water (WDW). The WDW, characterised by temperatures higher than 0 °C, occupies the majority of the water column above 1000 m except the upper 100-200 m, where Antarctic Surface Water (ASW) is found.

4.3. Relationship between zooplankton and acoustic backscatter

Based on first physical principles, sound is most efficiently scattered by objects of the size of the wavelength. The acoustic wavelength of our ADCPs, given by their nominal frequency of 75 kHz and a typical sound speed of 1500 m s^{-1} in seawater, is approximately 2 cm. This size class of zooplankton and nekton is represented by euphausiids, amphipods, pteropods, salps and small fish, for instance. Taking into account changes in body length during life cycles, other groups may also occur in the cm-range. However, animals with body sizes less than the wavelength by an order of magnitude can create strong backscatter signals too, if they are numerous enough to dominate the zooplankton assemblage as often reported from copepods, which represent the mm-size class (e.g. Pinot and Jansá, 2001). The backscatter strength of objects is also strongly influenced by their acoustic properties, in particular by their sound speed contrast to the environment, which depends mainly on their material composition. Organisms that contain for instance hard shells or gaseous enclosures in general scatter sound much stronger than gelatinous creatures. Concise overviews of the backscatter characteristics of certain zooplankton groups and their frequency-dependency can be found in Lavery et al. (2007) and in Warren and Wiebe (2008).

4.3.1. Species performing diel vertical migrations

Numerous net hauls of zooplankton were taken within the framework of the LAKRIS project, spread over the study area and during consecutive seasons and years. However, only a few hauls were made adjacent to the mooring positions and only at those certain instants of time, when the moorings were deployed and recovered. The number of pairs of ADCP profiles and net catches thus is much too small to relate particular acoustic signatures to certain zooplankton taxa. A further complication arises from the fact that most of the net hauls integrate vertically over depth ranges much too large to resolve the mostly narrow backscatter layers or have been taken just below the surface (Flores et al., 2014), where our acoustic records were too noisy for analysis. Consideration of the LAKRIS net catches in combination with other net catches collected from within the wider Weddell Sea however permits at least an idea of the different species that account for our acoustics-based observations.

Diel vertical migration is known from euphausiids but their behaviour may vary among species and in some cases between development stages of a species, being most common for adults (Nordhausen, 1994). DVM was observed in Antarctic krill, Euphausia superba (Godlewska, 1996; Siegel, 2005; Taki et al., 2005), Thysanoessa macrura (Nordhausen, 1994) and Euphausia crystallorophias (Everson, 1987). Taki et al. (2005) used Japanese fishery data from the Scotia Sea and analysed the average trawling depth of the Krill catches. Their results revealed a seasonal variation in the DVM behaviour of Euphausia superba, ranging from shallower residence depths during summer and early autumn, gradual deepening in autumn and maximum depths attained in winter. Euphausia superba and Thysanoessa macrura were the most abundant euphausiids in the eastern Weddell Sea during LAKRIS, while the other two euphausiid species that were collected, Euphausia frigida and Euphausia crystallorophias, were rare (Hunt et al., 2011; Siegel, 2012). Based on net catches made with an under-ice trawl, Flores et al. (2012) report for larval and postlarval krill (E. superba) a slightly higher daytime compared to nighttime near-surface abundance in summer but orders of magnitude higher nighttime than daytime accumulation near the surface in winter. Flores et al. (2014) showed that in autumn and winter, a deep mode of DVM was apparent from significant diel differences in Antarctic krill density of the epipelagic layer. Haraldsson and Siegel (2014) compared the length-frequency distribution and sex ratios of Thysanoessa macrura between shallow (0–200 m) and deep (0–2000 m) samples taken during winter. The authors found a distinct seasonal change in density and demography, together with the deeper samples; they concluded that Thysanoessa macrura performs a seasonal vertical migration in order to improve its survival during winter. By far the most abundant non-gelatinous zooplankton species in the size range most likely represented by our backscatter signal in the Lazarev Sea during all seasons were the euphausiids Thysanoessa macrura and Euphausia superba (Hunt et al., 2011; Flores et al., 2014). The seasonal and diurnal change in the vertical distribution of ADCP backscatter strength coincides in large parts with the known diel vertical migration behaviour of these species in the Lazarev Sea (Siegel, 2012; Haraldsson and Siegel, 2014; Flores et al., 2014). Seasonal and diel changes in the vertical distribution of euphausiids can therefore be considered to have most strongly influenced the observed patterns in ADCP backscatter.

DVM behaviour is also shown by other crustacean and noncrustacean zooplankton such as salps and pteropods (Nishikawa and Tsuda, 2001; Hunt et al., 2008), amphipods like *Themisto gaudichaudii* (Everson and Ward, 1980), and copepods (Atkinson et al., 1992) including small species (e.g. *Ctenocalanus citer*) and large species (e.g. *Rhincalanus gigas*). During the LAKRIS surveys, diel patterns of animal densities from net catches, with increased near-surface densities during the night consistent with DVM, were identified for *Clio pyramidata, Clione antarctica, Cyllopus lucasii, Hyperiella dilatata, Sagitta gazellae* and *Eukrohnia hamata* (Flores et al., 2011). Copepods also perform diel and seasonal vertical migrations, but are considered rather weak scatterers of sound (Stanton and Chu, 2000) at the frequency of 75 kHz used here.

The fish fauna sampled during the LAKRIS surveys was dominated by *Electrona antarctica* and *Notolepis coatsi* (Hunt et al., 2011). Both species possibly have also contributed to the acoustic backscattering. *Electrona antarctica* is known as a strong scatterer of sound as well as a diurnal vertical migrator with peak abundance at 0–300 m at night and 650–920 m during day (Greely et al., 1999). The same authors showed that its diet consists primarily of copepods, ostracods, and euphausiids, including *Euphausia superba*. In view of the observed backscattering levels, however, they should have occurred at low abundances (Benoit-Bird and Au, 2001).

4.3.2. Seasonal to interannual changes in zooplankton abundance

More than 100 species were found contributing to the macrozooplankton and micronekton community structure during LAKRIS (Hunt et al., 2011; Flores et al., 2014). Analysing catches from double oblique hauls with a Rectangular Midwater Trawl (RMT-8) in the upper 200 m of the water column Hunt et al. (2011) investigated the seasonal cycle of the macrozooplankton community. Copepods were not quantitatively sampled by the 4.5 mm RMT mesh size and thus were not considered in this analysis. Hunt et al. (2011) found that macrozooplankton densities did not differ significantly between seasons but that a shift towards a larger size structure occurred in the species assemblage between summer and autumn/winter. As a result, macrozooplankton biomass was higher in autumn and winter than in summer.

Compared to this net-based analysis of the macrozooplankton seasonal cycle, our acoustic estimates in terms of the water column mean volume backscatter strength, MVBS_{50-300m}, imply the contrary: highest zooplankton abundance in summer and lowest in winter. A very similar seasonal cycle as we derived from our moored 75 kHz ADCPs was obtained from a compilation of Drake Passage transits by a ship equipped with a 150 kHz ADCP sampling the depth layer 26–300 m, which indicated highest vertical mean backscatter strength in January and lowest in July (Chereskin and Tarling, 2007). Interesting to note is that even the amplitude of the seasonal cycle in Drake Passage south of the Polar Front is almost the same as in our observations in the eastern Weddell Sea, namely roughly 5 dB. One possible explanation for the similarity of our acoustic results with those of Chereskin and Tarling (2007) but their deviation from the RMT-based macrozooplankton analysis of Hunt et al. (2011) is congruence of the depth ranges considered by the two acoustic studies but their slight mismatch with the 0-200 m layer sampled by RMTs. Another possible explanation is that acoustic backscatter is sensitive to the bulk abundance of zooplankton, while net catches are rigorously size selective. Unresolved however, remains the question whether or not mesozooplankton such as copepods, in general too small to be adequately collected with the 4.5 mm RMT-8 mesh size, could have made a major contribution to the seasonal abundance amplitude implied by acoustics. While copepods are known to dominate the zooplankton assemblage in the Scotia and northern Weddell Sea (Ward et al., 2004) they are on the other hand considered

having only a weak acoustic target strength at 150 and particularly at 75 kHz (Stanton and Chu, 2000).

Most of the studies dedicated to determination of the relationship between zooplankton biomass and acoustic backscatter have found a linear increase of log-transformed dry weight of mixed zooplankton populations with MVBS, which is itself a logarithmic quantity (Fielding et al., 2004; Jiang et al., 2007; Warren and Wiebe, 2008). This relationship however was often hardly significant, and the regression slopes varied between studies. Nevertheless, if a zooplankton community is highly diverse as in our region (comprising more than 100 species not even including copepods) it may well be that acoustic backscatter information obtained from it, if averaged over many diurnal cycles and over a thick enough vertical layer, reveals most of the seasonal to interannual biomass variations. Using a tow package consisting of a multinet trawl apparatus and an ADCP mounted on the same frame directly in front of the net opening. Burd and Thomson (2012) have shown with nearly 200 sample pairs taken over large spatial scales and over 6 summers from deep sea zooplankton representing a large mix of faunal types and including animals with sizes ranging from copepods to small fish, that acoustic backscatter data explain 84% of the variance in total net biomass.

5. Summary and conclusion

Both methods of collecting in situ data for research into zooplankton dynamics, net catches and acoustic backscatter measurements, have their specific advantages and shortcomings. While single-frequency acoustic measurements hardly allow discrimination between zooplankton taxa and quantitative measurements of species biomass, they can yield long time series of the bulk zooplankton behaviour and abundance at high temporal resolution. Remoteness and harsh environmental conditions of the Southern Ocean notoriously have hampered the collection of scientific data, in particular from the winter season when access for ships to conduct net trawls is very restricted if not impossible. Acoustic backscatter time series records by moored instruments can provide valuable insights into the dynamics of zooplankton, and thus are useful in their own right. ADCPs have the additional advantage of measuring velocity in three dimensions, hence also vertical migration speeds of sound scattering organisms. Our 3-year long time series of MVBS and vertical velocity obtained from ADCPs moored at 64°S, 66.5°S and 69°S along the Greenwich Meridian from February 2005 until March 2008 provide a rare view into the diurnal to interannual variability of the abundance and behaviour of zooplankton in the Southern Ocean, and by combination with remote sensing and in situ data its responses to environmental forcing. Nocturnal diel vertical migration (DVM) of two distinct groups of migrators was found a highly persistent pattern at all three mooring locations and being closely related to the astronomical daylight cycles. And these observations correspond largely to the behaviour of the biomass dominants Euphausia superba and Thysanoessa macrura. However, while the DVM was symmetric around local noon, the annual modulation of the DVM was clearly asymmetric around winter solstice or summer solstice at all three mooring sites. Our data set confirms that the annual asymmetry, first noted by Cisewski et al. (2010), is a robust and recurring feature of the diel vertical migration behaviour in the eastern Weddell Sea. In contrast to several previous studies in other regions, DVM at our observation sites persisted throughout winter, even at the highest latitude during the polar night when the surface moreover was covered by sea ice. In all three years, a marked change in the migration behaviour occurred in late spring (late October/early November), when DVM ceased completely. The subsequent suspension of DVM after early November is likely caused by (1) increased availability of food in the surface mixed layer provided by the phytoplankton spring bloom, and (2) vanishing diel variation of the threat from visually oriented predators during the quasi-continuous illumination during the polar and subpolar summer. The highest migration speeds were recorded during the months February to April, i.e. from end of summer to early autumn, when the energy reserves of the zooplankter can reasonably be assumed filled to their annual maximum, and the slowest migration speeds were recorded during the months August to October, i.e. from end of winter to early spring, when the energy reserves are likely consumed after the food-poor winter period.

The mean annual cycle of the abundance of zooplankton in the water column, implied by the time series of normalised vertical mean backscatter strength, was characterised by a maximum in late summer and a gradual decrease during autumn and winter. It thus conforms to general wisdom about natural biological cycles in climatic zones with strong seasonal variation. The vernal increase followed upon the build-up of the phytoplankton spring bloom during the seasonal sea ice melt. However, the time series of water column mean backscatter strength from individual years and locations indicated interannual variations, which may exceed the seasonal amplitude of the mean, three years averaged, annual cycle. Comparison with satellite and in situ chlorophyll data revealed a considerable influence of the magnitude of the vernal pelagic phytoplankton bloom on the increase of the zooplankton abundance during the summer and its year-to-year variations. Without a phytoplankton spring bloom an increase of the zooplankton concentration may stay during the afflicted year. This leads to the conclusion that the annual growth of zooplankton, and in consequence the higher trophic levels in the Weddell Sea, depends mainly on the strength and areal extent of vernal pelagic phytoplankton bloom. However, while all the observed phytoplankton blooms occurred in shallow mixed layers left after the sea ice melt, not all melt water lenses supported phytoplankton blooms. Neither the sea ice coverage at the end of winter nor its vernal retreat and the associated mixed laver shallowing can be considered the primary determinator of the timing and magnitude of the seasonal phytoplankton bloom. As a complement to bottom up control we hypothesise top down control by zooplankton grazing on phytoplankton as a partial explanation for the observed interannual variations in the phytoplankton bloom magnitude. Our results suggest that a big phytoplankton bloom that developed during one summer can favour the growth of zooplankton on large horizontal scales to abundance levels that last for months and are still above the long-term average at the end of the following winter, and thus possibly high enough to graze down the phytoplankton in the subsequent spring, preventing the build-up of a bloom.

Acknowledgements

This work was co-funded by the German Federal Ministry of Education and Research (Bundesministerium für Bildung und Forschung, BMBF) through the joint project LAKRIS (Lazarev Sea Krill Study), which provided the salary of B.C. for three years and investment money for two ADCPs. We gratefully acknowledge the support provided by the captain, officers, and crew of the R/V *Polarstern.* Harry Leach, Harald Rohr and Timo Witte contributed substantially to the collection of the hydrographic data set. Axel Behrendt, Wolfgang Dierking and Hannelore Witte provided the upward looking sonar (ULS) data. The comments of an anonymous reviewer and of Hauke Flores in particular helped to perfect the paper.

References

- Ashjian, C.J., Smith, S.L., Flagg, C.N., Wilson, C., 1998. Patterns and occurrence of diel vertical migration of zooplankton biomass in the Mid-Atlantic Bight described by an acoustic Doppler current profiler. Continental Shelf Research 18, 831– 858.
- Ashjian, C.J., Campbell, R.G., Welch, H.E., Butler, M., Van Keuren, D., 2003. Annual cycle in abundance, distribution, and size in relation to hydrography of important copepod species in the western Arctic Ocean. Deep-Sea Research Part I 50, 1235–1261.
- Atkinson, A., Ward, P., Williams, R., Poulet, S.A., 1992. Diel vertical migration and feeding of copepods at an oceanic site near South Georgia. Marine Biology 113, 583–593.
- Atkinson, A., 1998. Life cycle strategies of epipelagic copepods in the Southern Ocean. Journal of Marine Systems 15, 289–311.
- Auerswald, L., Meyer, B., Teschke, M., Hagen, W., Kawaguchi, S., 2015. Physiological response of adult Antarctic krill, *Euphausia superba*, to long-term starvation. Polar Biology 38, 763–780.
- Bathmann, U.V., Makarov, R.R., Spiridonov, V.A., Rohardt, G., 1993. Winter distribution and overwintering strategies of the Antarctic copepod species *Calanoides acutus, Rhincalanus gigas* and *Calanus propinquus* (Crustacea, Calanoida) in the Weddell Sea. Polar Biology 13, 333–346.
- Bathmann, U. (Ed.), 2008. The Expedition ANTARKTIS-XXIII/6 of the Research Vessel "Polarstern" in 2006. Reports on Polar and Marine Research vol. 580, 1–168.
- Bathmann, U. (Ed.), 2010. The Expedition of the Research Vessel "Polarstern" to the Antarctic in 2007/2008 (ANT-XXIV/2). Reports on Polar and Marine Research vol. 604, 1–200.
- Behrendt, A., Fahrbach, E., Hoppema, M., Rohardt, G., Boebel, O., Klatt, O., Wisotzki, A., Witte, H., 2011. Variations of Winter Water properties and sea ice along the Greenwich meridian on decadal time scales. Deep-Sea Research Part II 58, 2524–2532.
- Behrendt, A., Dierking, W., Fahrbach, E., Witte, H., 2013. Sea ice draft in the Weddell Sea, measured by upward looking sonars. Earth System Science Data 5, 209– 226.
- Benoit-Bird, K.J., Au, W.W.L., 2001. Target strength measurements of Hawaiian mesopelagic boundary community animals. Journal of the Acoustical Society of America 110, 812–819.
- Benoit, D., Simard, Y., Gagne, J., Geoffroy, M., Fortier, L., 2010. From polar night to midnight sun: photoperiod, seal predation, and the diel vertical migration of polar cod (*Boreogadus saida*) under landfast ice in the Arctic Ocean. Polar Biology 33, 1505–1520.
- Berge, J., Cottier, F., Last, K.S., Varpe, Ø., Leu, E., Søreide, J., Eiane, K., Falk-Petersen, S., Willis, K., Nygård, H., Vogedes, D., Griffiths, C., Johnsen, G., Lorentzen, D., Brierley, A.S., 2009. Diel vertical migration of Arctic zooplankton during the polar night. Biology Letters 5, 69–72.
- Bersch, M., Becker, G.A., Frey, H., Koltermann, K.-P., 1992. Topographic effects of the Maud Rise on the stratification of the Weddell Gyre. Deep-Sea Research Part A 39, 303–331.
- Blachowiak-Samolyk, K., Kwasniewski, S., Richardson, K., Dmoch, K., Hansen, E., Hop, H., Falk-Petersen, S., Mouritsen, L.T., 2006. Arctic zooplankton do not perform diel vertical migration (DVM) during periods of midnight sun. Marine Ecology Progress Series 308, 101–116.
- Bogorov, B.G., 1946. Peculiarities of diurnal vertical migration of zooplankton in polar seas. Journal of Marine Research 6, 25–32.
- Brandt, A., Bathmann, U., Brix, S., Cisewski, B., Flores, H., Göcke, C., Janussen, D., Krägefsky, S., Kruse, S., Leach, H., Linse, K., Pakhomov, E., Peeken, I., Riehl, T., Sauter, E., Sachs, O., Schüller, M., Schrödl, M., Schwabe, E., Strass, V., van Franeker, J.A., Wilmsen, E., 2011. Maud Rise – a snapshot through the water column. Deep-Sea Research Part II 58, 1962–1982.
- Brierley, A.S., Saunders, R.A., Bone, D.G., Murphy, E.J., Enderlein, P., Conti, S.G., Demer, D.A., 2006. Use of moored acoustic instruments to measure short-term variability in abundance of Antarctic krill. Limnology and Oceanography: Methods 4, 18–29.
- Buchanan, C., Haney, J.F., 1980. Vertical migrations of zooplankton in the Arctic: a test of the environmental controls. In: Kerfoot, W.C. (Ed.), Evolution and Ecology of Zooplankton Communities. University Press of New England, Hanover, NH, pp. 69–79.
- Burd, B.J., Thomson, R.E., 2012. Estimating zooplankton biomass distribution in the water column near the Endeavour Segment of Juan de Fuca Ridge using acoustic backscatter and concurrently towed nets. Oceanography 25, 269–276. http://dx. doi.org/10.5670/oceanog.2012.25.
- Carmack, E.C., Foster, T.D., 1975. On the flow of water out of the Weddell Sea. Deep-Sea Research 22, 711–724.
- Chereskin, T.K., Tarling, G.A., 2007. Interannual to diurnal variability in the nearsurface scattering layer in Drake Passage. ICES Journal of Marine Science 64, 1617–1626.
- Cisewski, B., Strass, V.H., Prandke, H., 2005. Upper-ocean vertical mixing in the Antarctic Polar Front Zone. Deep-Sea Research Part II 52, 1087–1108.
- Cisewski, B., Strass, V.H., Rhein, M., Krägefsky, S., 2010. Seasonal variation of diel vertical migration of zooplankton from ADCP backscatter time series data in the Lazarev Sea, Antarctica. Deep-Sea Research Part I 57, 78–94.
- Cisewski, B., Strass, V.H., Leach, H., 2011. Circulation and transport of water masses in the Lazarev Sea, Antarctica, during summer and winter 2006. Deep-Sea Research Part I 58, 186–199.

- Clarke, A., Peck, L.S., 1991. The physiology of polar marine zooplankton. Polar Research 10, 355–369.
- Cohen, J.H., Forward, R.B., 2009. Zooplankton diel vertical migration a review of proximate control. Oceanography and Marine Biology an Annual Review 47, 77–110.
- Comiso, J., Gordon, A.L., 1987. Recurring polynyas over the Cosmonaut Sea and the Maud Rise. Journal of Geophysical Research Oceans 92, 2819–2833.
- Comiso, J., 1999, updated 2012. Bootstrap Sea Ice Concentrations from Nimbus-7 SMMR and DMSP SSM/I [01/01/2005–31/12/2008]. Digital Media. National Snow and Ice Data Center, Boulder, Colorado, USA.
- Connolley, W.M., 2005. Sea ice concentrations in the Weddell Sea: a comparison of SSM/I, ULS, and GCM data. Geophysical Research Letters 32, L07501. http://dx. doi.org/10.1029/2004GL021898.
- Cottier, F.R., Tarling, G.A., Wold, A., Falk-Petersen, S., 2006. Unsynchronized and synchronized vertical migration of zooplankton in a high Arctic fjord. Limnology and Oceanography 51, 2586–2599.
- Deines, K.L., 1999. Backscatter estimation using broadband acoustic Doppler current profiles. IEEE, 249–253.
- de Steur, L., Holland, D.M., Muench, R.D., McPhee, M.G., 2007. The warm-water "Halo" around Maud Rise: properties, dynamics and Impact. Deep-Sea Research Part I 54, 871–896.
- Eiane, K., Aknes, D.L., Ohman, M.D., 1998. Advection and zooplankton fitness. Sarsia 83, 87–93.
- Everson, I., Ward, P., 1980. Aspects of Scotia Sea zooplankton. Biological Journal of the Linnean Society 14, 93–101.
- Everson, I., 1987. Some aspects of the small scale distribution of *E. crystallorophias*. Polar Biology 8, 9–15.
- Fach, B., Schmidt, C., Auerswald, L., Hayden, A., Herrmann, R., Hohn, S., Krägefsky, S., Meyer, B., 2007. Distribution of chlorophyll a in the Lazarev Sea. Reports on Polar and Marine Research vol. 568, 56–57.
- Fahrbach, E., Hoppema, M., Rohardt, G., Schröder, M., Wisotzki, A., 2004. Decadalscale variations of water mass properties in the deep Weddell Sea. Ocean Dynamics 54, 77–91.
- Falk-Petersen, S., Leu, E., Berge, J., Kwasniewski, S., Nygård, H., Røstad, A., Keskinen, E., Thormar, J., von Quillfeldt, C., Wold, A., Gulliksen, B., 2008. Vertical migration in high Arctic waters during autumn 2004. Deep-Sea Research Part II 55, 2275– 2284.
- Fielding, S., Griffiths, G., Roe, H.S.J., 2004. The biological validation of ADCP acoustic backscatter through direct comparison with net samples and model predictions based on acoustic-scattering models. ICES Journal of Marine Science 61, 184– 200.
- Firing, E., 1991. Acoustic Doppler Current Profiling Measurements and Navigation, WOCE Hydrographic Program Office Report. WHPO 91-9, WOCE Report. 68/91, 24 pp.
- Fischer, J., Visbeck, M., 1993. Seasonal variation of the daily zooplankton migration in the Greenland Sea. Deep-Sea Research Part I 40, 1547–1557.
- Flagg, C.N., Smith, S.L., 1989. On the use of the acoustic Doppler current profiler to measure zooplankton abundance. Deep-Sea Research Part A 36, 455–474.
- Flagg, C.N., Wirick, C.D., Smith, S.L., 1994. The interaction of phytoplankton, zooplankton and currents from 15 months of continuous data in the Mid-Atlantic Bight. Deep-Sea Research Part II 41, 411–435.
- Flores, H., van Franeker, J.A., Cisewski, B., Leach, H., Van de Putte, A.P., Meesters, E.H. W.G., Bathmann, U., Wolff, W.J., 2011. Macrofauna under sea ice and in the open surface layer of the Lazarev Sea, Southern Ocean. Deep-Sea Research Part II 58, 1948–1961.
- Flores, H., van Franeker, J.A., Siegel, V., Haraldsson, M., Strass, V., Meesters, E.H., Bathmann, U., Wolff, W.J., 2012. The association of Antarctic krill *Euphausia superba* with the under-ice habitat. PLoS ONE 7, e31775. http://dx.doi.org/ 10.1371/journal.pone.0031775.
- Flores, H., Hunt, B.P.V., Kruse, S., Pakhomov, E.A., Siegel, V., van Franeker, J.A., Strass, V., Van de Putte, A.P., Meesters, E.H.W.G., Bathmann, U., 2014. Seasonal changes in the vertical distribution and community structure of Antarctic macrozooplankton and micronekton. Deep-Sea Research Part I 84, 127–141.
- Fofonoff, N.P., Millard, R.C., 1983. Algorithms for the computation of fundamental properties of seawater. UNESCO Technical Papers in Marine Science 44, 1–53.
- Fortier, M., Fortier, L., Hattori, H., Saito, H., Legendre, L., 2001. Visual predators and the diel vertical migration of copepods under Arctic sea ice during the midnight sun. Journal of Plankton Research 23, 1263–1278.
- Forward, R.B., 1988. Diel vertical migration: zooplankton photobiology and behaviour. Oceanography and Marine Biology Annual Review 26, 361–393.
- Francois, R.E., Garrison, G.R., 1982. Sound absorption based upon ocean measurements. Part II: boric acid contribution and equation for total absorption. Journal of the Acoustical Society of America 72, 1879–1890.
- Gloersen, P., Campbell, W.J., 1991. Recent variations in Arctic and Antarctic sea-ice covers. Nature 352, 33–36.
- Godlewska, M., 1996. Vertical migrations of krill (*Euphausia superba* Dana). Polish Archiwum Hydrobiologii 43 (1), 9–63.
- Gordon, A.L., Huber, B.A., 1995. Warm Weddell deep water west of Maud Rise. Journal of Geophysical Research Oceans 100, 13747–13753.
- Greely, T.M., Gartner, J.V., Torres, J.J., 1999. Age and growth of *Electrona antarctica* (Pisces: Myctophidae), the dominant mesopelagic fish of the Southern Ocean. Marine Biology 133, 145–158.
- Hagen, W., Schnack-Schiel, S.B., 1996. Seasonal lipid dynamics in dominant Antarctic copepods: energy for overwintering or reproduction? Deep Sea Research Part I 43, 139–158.

- Hagen, W., Kattner, G., Terbrüggen, A., Van Vleet, E.S., 2001. Lipid metabolism of the Antarctic krill *Euphausia superba* and its ecological implications. Marine Biology 139, 95–104.
- Haney, J.F., 1988. Diel patterns of zooplankton behaviour. Bulletin of Marine Science 43, 583–603.
- Haraldsson, M., Siegel, V., 2014. Seasonal distribution and life history of *Thysanoessa* macrura (Euphausiacea, Crustacea) in high latitude waters of the Lazarev Sea, Antarctica. Marine Ecology Progress Series 495, 105–118.
- Hardy, A.C., Gunther, E.R., 1935. The plankton of South Georgia whaling grounds and adjacent waters 1926–1927. Discovery Report 11, 1–456.
- Hays, G.C., 1995. Ontogenetic and seasonal variation in the diel vertical migration of the copepods *Metridia lucens* and *Metridia longa*. Limnology and Oceanography 40, 1461–1465.
- Herrmann, S., 2009. Das Mikrophytoplankton der Lazarewsee im antarktischen Winter Verteilung, Abundanz und Artenzusammensetzung. Diploma Thesis. Faculty of Biology and Chemistry, University of Bremen, Bremen.
- Heywood, K.J., Scrope-Howe, S., Barton, E.D., 1991. Estimation of zooplankton abundance from shipborne ADCP backscatter. Deep-Sea Research Part A 38, 677–691.
- Heywood, K.J., 1996. Diel vertical migration of zooplankton in the northeast Atlantic. Journal of Plankton Research 18, 163–184.
- Hunt, B.P.V., Pakhomov, E.A., Hosie, G.W., Siegel, V., Ward, P., Bernard, K., 2008. Pteropods in Southern Ocean ecosystems. Progress in Oceanography 78, 193– 221.
- Hunt, B.P.V., Pakhomov, E.A., Siegel, V., Strass, V., Cisewski, B., Bathmann, U., 2011. The seasonal cycle of the Lazarev Sea macrozooplankton community and a potential shift to top-down trophic control in winter. Deep-Sea Research Part II 58, 1662–1676.
- Jacobs, S.S., 1991. On the nature and significance of the Antarctic Slope Front. Marine Chemistry 35, 9–24.
- Jiang, S.N., Dickey, T.D., Steinberg, D.K., Madin, L.P., 2007. Temporal variability of zooplankton biomass from ADCP backscatter time series data at the Bermuda Testbed Mooring site. Deep-Sea Research Part I 54, 608–636.
- Kattner, G., Hagen, W., 2009. Lipids in marine copepods: latitudinal characteristics and perspective to global warming. In: Arts, M.T., Brett, M.T., Kainz, M. (Eds.), Lipids in Aquatic Ecosystems. Springer, New York, NY, pp. 257–280.
- Kawaguchi, K., Ishikawa, S., Matsuda, O., 1986. The overwintering strategy of Antarctic krill (*Euphausia superba* Dana) under the coastal fast ice off the Ongul Islands in Lützow-Holm Bay, Antarctica. Memoirs of National Institute of Polar Research 44, 67–85.
- Krägefsky, S., 2008. On the Copepod Response to Iron-induced Phytoplankton Blooms in the Southern Ocean Ph.D. Thesis. University of Bremen, Germany, pp. 1–239.
- Krägefsky, S., Bathmann, U., Strass, V., Wolf-Gladrow, D., 2009. Response of small copepods to an iron-induced phytoplankton bloom – a model to address the mechanisms of aggregation. Marine Ecology Progress Series 374, 181–198.
- Lampert, W., 1989. The adaptive significance of diel vertical migration of zooplankton. Functional Ecology 3, 21–27.
- Lavery, A.C., Wiebe, P.H., Stanton, T.K., Lawson, G.L., Benfield, M.C., Copley, N., 2007. Determining dominant scatterers of sound in mixed zooplankton populations. Journal of the Acoustical Society of America 122, 3304–3326. http://dx.doi.org/ 10.1121/1.2793613.
- Leach, H., Strass, V.H., Cisewski, B., 2011. Modification by lateral mixing of the warm deep water entering the Weddell Sea in the Maud Rise region. Ocean Dynamics 61, 51–68. http://dx.doi.org/10.1007/s10236-010-0342-y.
- Longhurst, A.R., Harrison, W.G., 1988. Vertical nitrogen flux from the oceanic photic zone by diel migrant zooplankton and nekton. Deep-Sea Research Part A 35, 881–889.
- Luo, J., Ortner, P.B., Forcucci, D., Cummings, S.R., 2000. Diel vertical migration of zooplankton and mesopelagic fish in the Arabian Sea. Deep-Sea Research Part II 47, 1451–1473.
- Manuel, J.L., O'Dor, R.K., 1997. Vertical migration for horizontal transport while avoiding predators: I. A tidal/diel model. Journal of Plankton Research 19, 1929– 1947.
- Maritorena, S., Siegel, D.A., 2005. Consistent merging of satellite ocean color data sets using a bio-optical model. Remote Sensing of Environment 94, 429–440.
- Massom, R.A., Stammerjohn, S.E., 2010. Antarctic sea ice change and variability physical and ecological implications. Polar Science 4, 149–186.
- Meyer, B., Auerswald, L., Siegel, V., Spahic, S., Pape, C., Fach, B.A., Teschke, M., Lopata, A.L., Fuentes, V., 2010. Seasonal variation in body composition, metabolic activity, feeding, and growth of adult krill *Euphausia superba* in the Lazarev Sea. Marine Ecological Progress Series 398, 1–18.
- Meyer, B., 2012. The overwintering of Antarctic krill, *Euphausia superba*, from an ecophysiological perspective. Polar Biology 35, 15–37. http://dx.doi.org/ 10.1007/s00300-011-1120-0.
- Muench, R.D., Morison, J.H., Padman, L., Martinson, D., Schlosser, P., Huber, B., Hohmann, R., 2001. Maud Rise revisited. Journal of Geophysical Research Oceans 106, 2423–2440.
- Nishikawa, J., Tsuda, A., 2001. Diel vertical migration of the tunicate *Salpa thompsoni* in the Southern Ocean during summer. Polar Biology 24, 299–302.
- Nordhausen, W., 1994. Distribution and diel vertical migration of the euphausiid *Thysanoessa macrura* in Gerlache Strait, Antarctica. Polar Biology 14, 219–229.
- Ou, H.W., 1991. Some effects of a seamount on oceanic flows. Journal of Physical Oceanography 21, 1835–1845.
- Padman, L., Fricker, H.A., Coleman, R., Howard, S., Erofeeva, L., 2002. A new tide model for the Antarctic ice shelves and seas. Annals of Glaciology 34, 247–254.

- Park, Y.-H., Charriaud, E., Fieux, M., 1998. Thermohaline structure of the Antarctic Surface Water/Winter Water in the Indian sector of the Southern Ocean. Journal of Marine Systems 17, 5–23.
- Pinot, J.M., Jansa, J., 2001. Time variability of acoustic backscatter from zooplankton in the Ibiza Channel (western Mediterranean). Deep-Sea Research Part I 48, 1651–1670.
- Plueddemann, A.J., Pinkel, R., 1989. Characterization of the patterns of diel migration using a Doppler sonar. Deep-Sea Research Part A 36, 509–530.
- Quetin, L.B., Ross, R.M., 2009. Life under Antarctic pack ice: a krill perspective. In: Krupnik, I., Lang, M.A., Miller, E.S. (Eds.), Smithsonian at the Poles, Contribution International Polar Year Science. Smithsonian Institute, Washington, DC, pp. 285–298.
- Rabindranath, A., Daase, M., Falk-Petersen, S., Wold, A., Wallace, M.I., Berge, J., Brierley, A.S., 2010. Seasonal and diel vertical migration of zooplankton in the High Arctic during autumn midnight sun of 2008. Marine Biodiversity 41, 356– 382. http://dx.doi.org/10.1007/s12526-010-0067-7.
- Radenac, M.-H., Plimpton, P.E., Lebourges-Dhaussy, A., Commien, L., McPhaden, M.J., 2010. Impact of environmental forcing on the acoustic backscattering strength in the equatorial Pacific: diurnal, lunar, intraseasonal and interannual variability. Deep-Sea Research Part I 57, 1314–1328.
- Reda, I., Andreas, A., 2004. Solar position algorithm for solar radiation applications. Solar Energy 76, 577–589.
- Ringelberg, J., 1995. Changes in light intensity and diel vertical migration: a comparison of marine and freshwater environments. Journal of the Marine Biology Association of the United Kingdom 75, 15–25.
- Robertson, R., Padman, L., Egbert, G.D., 1988. Tides in the Weddell Sea. Ocean, Ice, and Atmosphere: Interactions at the Antarctic Continental Margin Antarctic Research Series 75, 341–369.
- Roe, H.S.J., 1974. Observations on the diurnal vertical migrations of an oceanic animal community. Marine Biology 28, 99–113.
- Schenke, H.W., Dijksta, S., Niederjasper, F., Schöne, T., Hinze, H., Hoppman, B., 1998. The new bathymetric charts of the Weddell Sea: AWI BCWS. In: Jacobs, S.S., Weiss, R.F. (Eds.), Ocean, Ice, and Atmosphere: Interactions at the Antarctic Continental Margin, Antarctic Research Series, vol. 75. AGU, Washington, DC, pp. 371–380. http://dx.doi.org/10.1029/AR075p0371.
- Siegel, V., 2005. Distribution and population dynamics of *Euphausia superba*: summary of recent findings. Polar Biology 29, 1–22.
- Siegel, V., 2012. Krill stocks in high latitudes of the Antarctic Lazarev Sea: seasonal and interannual variation in distribution, abundance and demography. Polar Biology 35, 1151–1177.
- Smetacek, V., Klaas, C., Strass, V., Assmy, P., Montresor, M., Cisewski, B., Savoye, N., Webb, A., d'Ovidio, F., Arrieta, J.M., Bathmann, U., Bellerby, R., Berg, G.M., Croot, P.L., Gonzalez, S., Henjes, J., Herndl, G.J., Hoffmann, L.J., Leach, H., Losch, M., Mills, M.M., Neill, C., Peeken, I., Roettgers, R., Sachs, O., Sauter, E., Schmidt, M., Schwarz, J.N., Terbrüggen, A., Wolf-Gladrow, D., 2012. Deep carbon export from a Southern Ocean iron-fertilized diatom bloom. Nature 487 (7407), 313– 319.
- Schmidt, K., Atkinson, A., Pond, D.W., Ireland, L.C., 2014. Feeding and overwintering of Antarctic krill across its major habitats: the role of sea ice cover, water depth, and phytoplankton abundance. Limnology and Oceanography 59, 17–36.
- Smith, W.O., Nelson, D.M., 1985. Phytoplankton bloom produced by a receding ice edge in the Ross Sea: spatial coherence with the density field. Science 227 (4683), 163–166.
- Stanton, T.K., Chu, D., 2000. Review and recommendations for the modelling of acoustic scattering by fluid-like elongated zooplankton: euphausiids and copepods. ICES Journal of Marine Science 57, 793–807.
- Steinberg, D.K., Carlson, C.A., Bates, N.R., Goldthwait, S.A., Madin, L.P., Michaels, A.F., 2000. Zooplankton vertical migration and the active transport of dissolved organic and inorganic carbon in the Sargasso Sea. Deep-Sea Research Part I 47, 137–158.
- Strass, V.H., 1998. Measuring sea ice draft and coverage with moored upward looking sonars. Deep-Sea Research Part I 45, 795–818.
- Strass, V. (Ed.), 2007. The expedition ANTARKTIS-XXIII/2 of the Research Vessel "Polarstern" in 2005/2006. Reports on Polar and Marine Research vol. 568, 1–138.
- Taki, K., Hayashi, T., Naganobu, M., 2005. Characteristics of seasonal variation in diurnal vertical migration and aggregation of Antarctic krill (*Euphausia superba*) in the Scotia Sea, using Japanese fishery data. CCAMLR Science 12, 163–172.
- Tarling, G.A., Matthews, J.B.L., David, P., Guerin, O., Buchholz, F., 2001. The swarm dynamics of northern krill (*Meganyctiphanes norvegica*) and pteropods (*Cavolinia inflexa*) during vertical migration in the Ligurian Sea observed by an acoustic Doppler current profiler. Deep-Sea Research Part I 48, 1671– 1686.
- Taylor, M.H., Losch, M., Bracher, A., 2013. On the drivers of phytoplankton blooms in the Antarctic marginal ice zone: a modeling approach. Journal of Geophysical Research Oceans 118, 63–75.
- Teschke, M., Wendt, S., Kawaguchi, S., Kramer, A., Meyer, B., 2011. A circadian clock in Antarctic Krill: an endogenous timing system governs metabolic output rhythms in the euphausid species *Euphausia superba*. PLoS ONE 6, e26090. http://dx.doi.org/10.1371/journal.pone.0026090.
- Torres, J.J., Donnelly, J., Hopkins, T.L., Lancraft, T.M., Aarset, A.V., Ainley, D.G., 1994. Proximate composition and overwintering strategies of Antarctic micronektonic Crustacea. Marine Ecology Progress Series 113, 221–232.
- Turner, J.T., 2015. Zooplankton fecal pellets, marine snow, phytodetritus and the ocean's biological pump. Progress in Oceanography 130, 205–248. http://dx.doi. org/10.1016/j.pocean.2014.08.005.

- van Haren, H., Compton, T.J., 2013. Diel vertical migration in deep sea plankton is finely tuned to latitudinal and seasonal day length. PLoS ONE 8, e64435. http:// dx.doi.org/10.1371/journal.pone.0064435.
- Vélez-Belchí, P., Allen, J.T., Strass, V., 2002. A new way to look at mesoscale zooplankton distributions: an application at the Antarctic Polar Front. Deep-Sea Research Part II 49, 3917–3929.
- Wallace, M.I., Cottier, F.R., Berge, J., Tarling, G.A., Griffiths, C., Brierley, A.S., 2010.
 Comparison of zooplankton vertical migration in an ice-free and seasonally icecovered Arctic fjord: an insight into the influence of sea ice cover on zooplankton behaviour. Limnology and Oceanography 55, 831–845.
 Wallace, M.I., Cottier, F.R., Brierley, A.S., Tarling, G.A., 2013. Modelling the influence
- Wallace, M.I., Cottier, F.R., Brierley, A.S., Tarling, G.A., 2013. Modelling the influence of copepod behaviour on faecel pellet export at high latitudes. Polar Biology 36, 579–592.
- Ward, P., Grant, S., Brandon, M., Siegel, V., Sushin, V., Loeb, V., Griffiths, H., 2004. Mesozooplankton community structure in the Scotia Sea during the CCAMLR 2000 survey: January–February 2000. Deep-Sea Research Part II 51, 1351–1367.
- Warren, J.D., Wiebe, P.H., 2008. Accounting for biological and physical sources of acoustic backscatter improves estimates of zooplankton biomass. Canadian Journal of Fisheries and Aquatic Sciences 65 (7), 1321–1333.
- Wroblewski, J.S., 1982. Interaction of currents and vertical migration in maintaining Calanus marshallae in the Oregon upwelling zone – a simulation. Deep-Sea Research Part A 29, 665–668.
- Zaret, T.M., Suffern, J.S., 1976. Vertical migration in zooplankton as a predator avoidance mechanism. Limnology and Oceanography 21, 804–813.
- Zwally, H.J., Parkinson, C.L., Comiso, J.C., 1983. Variability of Antarctic sea ice and changes in carbon dioxide. Science 220 (4601), 1005–1012.